

# MODULAR PROPERTIES OF RIBBON ABELIAN CATEGORIES

VOLODIMIR LYUBASHENKO

**Abstract.** A category  $N$  of labeled (oriented) trivalent graphs (nets) or ribbon graphs is extended by new generators called fusing, braiding, twist and switch with relations which can be called Moore–Seiberg relations. A functor to  $N$  is constructed from the category  $Surf$  of oriented surfaces with labeled boundary and their homeomorphisms. Given an (eventually non-semisimple)  $k$ -linear abelian ribbon braided category  $\mathcal{C}$  with some finiteness conditions we construct a functor from a central extension of  $N$  with the set of labels  $\text{Ob}\mathcal{C}$  to  $k$ -vector spaces. Composing the functors we get a modular functor from a central extension of  $Surf$  to  $k$ -vector spaces.

1991 Mathematics Subject Classification: 18B30, 18D10, 57N05

Moore and Seiberg’s study [16] on conformal field theory was continued and developed by Walker [23] from the topological point of view. The first systematic study in this direction of the example of  $\widehat{\mathfrak{sl}}(2)$  was made by Kohno [5, 6]. A different topological approach was proposed by Reshetikhin and Turaev [18] (see also [21]). The aim of this article is to present a categorical point of view on the subject. We use freely notations and results from the previous papers [9, 10].

We consider a category of surfaces  $\mathcal{S}$  labeled by a set  $\mathcal{C}$ , which Grothendieck calls the Teichmüller’s tower. Its subcategory  $Surf$  consists of labeled surfaces and isotopy classes of their homeomorphisms. Its central extension is denoted  $ESurf$ . We give also a definition of a modular functor.

We show that any ribbon abelian category  $\mathcal{C}$ , satisfying the axioms of modularity [10] yields a modular functor  $Z_{\mathcal{C}} : ESurf \rightarrow k\text{-vect}$ . Thus, such category deserves to be called *modular*. Precisely, modularity means the following:  $\mathcal{C}$  is a noetherian abelian  $k$ -linear ribbon tensor category with finite dimensional  $k$ -vector spaces  $\text{Hom}_{\mathcal{C}}(A, B)$ .

---

*Date:*

The research was supported in part by the SERC research grant GR/G 42976.  
The detailed version of this paper has been submitted for publication elsewhere.

In a cocompletion of  $\mathcal{C}$  there exists a Hopf algebra  $F = \int^{X \in \mathcal{C}} X \otimes X^\vee$ , an automorphism  $T : F \rightarrow F$  and a Hopf pairing  $\omega : F \otimes F \rightarrow I$  (see [10]) with the kernel  $\text{Ker } \omega \subset F$ . We require that

- (M1)  $\mathbf{f} \stackrel{\text{def}}{=} F / \text{Ker } \omega$  is an object of  $\mathcal{C}$  (“ $\mathbf{f}$  is finite dimensional”);
- (M2)  $T(\text{Ker } \omega) \subset \text{Ker } \omega$ .

Vice versa, with some assumptions of representability any modular functor  $Z : ESurf \rightarrow k\text{-vect}$  induce on  $\mathcal{C}$  the structure of a ribbon category and the functor  $Z$  is isomorphic to  $Z_{\mathcal{C}}$ . So, up to some extent, our conditions are not only sufficient, but also necessary.

In the case when  $\mathcal{C}$  is a semisimple abelian category with finite number of simple objects and  $\text{Ker } \omega = 0$ , obtained results generalize those of Moore and Seiberg [16]. Practically, in this case the obtained functor is the same as constructed by Walker [23] (in dimension two) and probably coincides with the one of Reshetikhin and Turaev [18]. A new feature is a possibility to work with non-semisimple categories.

**Acknowledgements .** I wish to thank V.G. Drinfeld, A. Joyal, T. Kohno, S. Majid, Y. Soibelman, R. Street for multiple helpful discussions. I am grateful to L. Breen for his kind hospitality and opportunity to work in Université Paris-Nord, where some results of this paper were obtained. More results were obtained, when the author visited C. De Concini at Scuola Normale Superiore and I express my deep gratitude to him.

## 1. Introduction

In Section 2 a category of labeled surfaces  $\mathcal{S}$  and isotopy classes of their glueing maps is considered. It encompasses the usual category of surfaces  $Surf$  with isotopy classes of homeomorphisms as morphisms. This is a topological counterpart of Teichmüller’s tower, considered by Grothendieck [3]. He conjectured that the Teichmüller’s tower can be described by few essential relations in small genera. These relations were written by Moore and Seiberg [16].

They are essential relations of the category of ribbon graphs  $RG$  defined in Section 2.2. Ribbon graphs are surfaces with distinguished intervals at the boundary. The category has infinite number of generators combined into finite number of classes: topological morphisms (homeomorphisms and glueings of ribbon graphs), braiding, twists and switches. Also it has infinite number of relations, which are relatively trivial and can be considered as commutation relations or as identification of generators.

There is a functor  $dupl : RG \rightarrow \mathcal{S}$ . We construct in Section 3 a functor  $Surf \rightarrow RG$  using the presentations of the mapping class group  $M_{g,0}$  by Wajnryb [22] and of the braid group of a closed surface by Scott [19]. The composition  $Surf \rightarrow RG \rightarrow \mathcal{S}$  is the natural inclusion. The conjecture is that the functor  $dupl$  is an equivalence between  $\mathcal{S}$  and  $RG$ .


We describe another category which objects are 1-complexes (graphs) called nets and morphisms besides maps of graphs include morphisms of insertion or deletion of a vertex, fusing, braiding, twists and switches. This category comes in two versions—unoriented  $N$  and oriented  $ON$  (Sections 4–6). Both are equivalent to the category of ribbon graphs  $RG$ .

In Section 7 we study central extensions of the last category (in any form) and define a universal extension  $EN$ .

Given an abelian ribbon tensor category  $\mathcal{C}$  with some finiteness properties we define a functor on a category  $EN$  labeled by objects of  $\mathcal{C}$  in Section 9. Composing the functors we get a representation of the central extension of the category  $Surf$ . This is a direct generalization of Moore and Seiberg’s results [16].

Examples of such categories  $\mathcal{C}$  can be found in [12]. Besides familiar semisimple abelian categories with finite number of simple objects, there are other, non-semisimple examples, such as the category of representations of a quantum group at a root of unity.

All the relevant categories and functors make part of the following diagram



Proofs of the obtained results will be published elsewhere [11].

## 2. Surfaces

**2.1. Labeled surfaces.** Let  $\mathcal{C} = \{A, B, C, \dots\}$  be a set of labels with an involution  $\cdot^\sim : \mathcal{C} \rightarrow \mathcal{C}$ . By a *labeled surface* we shall understand

the following: a compact oriented surface  $\Sigma$  with a boundary, with a labelling of boundary circles  $L : \pi_0(\partial\Sigma) \rightarrow \mathcal{C}$ ,  $i \mapsto A_i$  and with a chosen point  $x_i$  on  $i^{\text{th}}$  boundary circle, i.e. a section  $x : \pi_0(\partial\Sigma) \rightarrow \partial\Sigma$  of the projection  $\partial\Sigma \rightarrow \pi_0(\partial\Sigma)$  is fixed. So we write  $\partial\Sigma = \coprod_i \underline{C}_i$ , where a circle is described as  $\underline{C}_i = (C_i, A_i, x_i)$ .

**Definition 2.1.** Let  $\Sigma, \tilde{\Sigma}$  be two labeled surfaces. A continuous surjective mapping  $f : \Sigma \rightarrow \tilde{\Sigma}$  is called *glueing*, if its restriction to interior part is an orientation preserving homeomorphism  $\text{Int } \Sigma \rightarrow f(\text{Int } \Sigma)$  and  $\tilde{\Sigma}$  is a quotient space of  $\Sigma$  with respect to the following equivalence relation  $\sim$  in  $\partial\Sigma$ . We choose in  $\pi_0(\partial\Sigma)$  some mutually disjoint pairs  $(C'_j, A_j, x'_j)$  and  $(C''_j, A_j, x''_j)$  and homeomorphisms  $\phi_j : C'_j \rightarrow C''_j$  such that  $\phi_j(x'_j) = x''_j$ . If  $\phi_j(x') = x''$ , we write  $x' \sim x''$ . For any boundary circle  $(C_i, A_i, x_i)$  in  $\tilde{\Sigma}$  its preimage must have the same labels  $(f^{-1}(C_i), A_i, f^{-1}(x_i))$ .

Note that a particular case of a glueing is an orientation, labelling and distinguished points preserving homeomorphism. Since a composition of glueings is again a glueing, they form a category.

**Definition 2.2.** Consider a category  $\mathcal{S} = \mathcal{S}_{\mathcal{C}}$  which objects are labeled surfaces and morphisms are isotopy classes of glueings (an isotopy here is a continuous family of glueings). The category  $\mathcal{S}$  is a symmetric monoidal category with disjoint union as a monoidal product.

**Proposition 2.1.** *The equality  $gf_1 = gf_2$  of morphisms in the category  $\mathcal{S}$  implies  $f_1 = f_2$  (we shall say that the category  $\mathcal{S}$  has left cancellations property).*

*Proof.* Glueings are surjective maps. □

## 2.2. Ribbon graphs.

**Definition 2.3.** A ribbon graph is an oriented surface  $X$  such that its each component has a non-empty boundary, equipped with a subset  $B \subset \partial X$  homeomorphic to finite disjoint sum of closed intervals, with a labelling  $L : \pi_0(B) \rightarrow \mathcal{C}$ , and with a chosen point in the interior of each component of  $B$ , i.e. with a section  $x : \pi_0(B) \rightarrow \text{Int } B$  of the projection  $B \rightarrow \pi_0(B)$ .

**Definition 2.4.** Let  $\mathcal{RG}$  be a category which objects are ribbon graphs and morphisms are isotopy classes of glueings of ribbon graphs. A *glueing*  $f : (X_1, B_1, L_1) \rightarrow (X_2, B_2, L_2)$  is a continuous surjective orientation preserving mapping  $f : X_1 \rightarrow X_2$ , such that the preimage of each point consists of one or two points and the induced relation  $\sim$  in  $X_1$

$(x, y \in X_1$  are equivalent if  $f(x) = f(y))$  reduces to pairwise identification of some components of  $B_1$  with other components having dual labels. Distinguished points must be pairwise identified or mapped to distinguished points, preimage  $f^{-1}(B_2)$  must be a union of components of  $B_1$ , and homeomorphism  $f : f^{-1}(B_2) \rightarrow B_2$  must preserve labeling. The category  $\mathcal{RG}$  is a symmetric monoidal category with respect to disjoint union.

The category  $\mathcal{RG}$  has left cancellation property.

There is a functor  $dupl : \mathcal{RG} \rightarrow \mathcal{S}$  called duplication. Having a ribbon graph  $X$ , construct a surface  $\Sigma = X \cup \bar{X}$ , where  $\bar{X}$  is a second copy of  $X$  with reversed orientation and  $\partial X - \text{Int } B$  is identified with  $\partial \bar{X} - \text{Int } \bar{B}$  via “identity map”. Boundary of  $\Sigma$  is the suspension of  $B$  and it obtains labeling from the labeling of  $B$ . The chosen points in  $B$  become chosen points in  $\partial \Sigma$ . To each glueing  $f : X \rightarrow Y$  of ribbon graphs corresponds the glueing  $f \cup \bar{f} : X \cup \bar{X} \rightarrow Y \cup \bar{Y}$  of surfaces.

The duplication functor is injective on morphisms and essentially surjective on objects. Duplication of the following ribbon graphs (double lines mark the subset  $B \subset \partial X$ )

$$D_0 = \bigcirc, \quad D_1 = \overline{\text{A}}, \quad D_2 = \begin{array}{|c|} \hline \text{A} \\ \hline \text{E} \\ \hline \end{array}, \quad D_3 = \begin{array}{c} \text{C} \\ \text{A} \quad \text{E} \end{array} \quad (2.1)$$

$$A_0 = \text{torus}, \quad A_1 = \text{torus with one hole} \quad (2.2)$$

gives sphere, disk, annulus, pants, torus and torus with one hole correspondingly.

The category  $\mathcal{S}$  has more morphisms than  $\mathcal{RG}$ . For instance, the following are automorphisms of annulus, pants (sphere  $S^2 = \bar{\mathbb{C}}$  with 3

holes) and torus with one hole

$$\begin{array}{ccc}
 R : & \begin{array}{c} \text{Diagram 1: A square with rounded corners, labeled } A \text{ at the top. Inside is a circle labeled } B. \text{ A vertical line segment connects } A \text{ to } B. \end{array} & \longrightarrow & \begin{array}{c} \text{Diagram 2: A square with rounded corners, labeled } A \text{ at the top. Inside is a circle labeled } B. \text{ A line segment connects } A \text{ to } B, \text{ and a curved line segment connects } B \text{ back to } A, \text{ forming a loop.} \end{array} \\
 \\
 \omega = {}_C\omega_{AE} : & \begin{array}{c} \text{Diagram 3: A tree graph with root } C \text{ at the top, and children } A \text{ and } E \text{ at the bottom.} \end{array} & \longrightarrow & \begin{array}{c} \text{Diagram 4: A graph with root } C \text{ at the top, and children } E \text{ and } A \text{ at the bottom. A curved line segment connects } E \text{ and } A, \text{ forming a loop.} \end{array} \quad (2.3)
 \end{array}$$

$$\begin{array}{ccc}
 S : & \begin{array}{c} \text{Diagram 5: A square divided into four quadrants by a horizontal and vertical line. A circle is in the bottom-left quadrant, connected to the horizontal line by a vertical segment.} \end{array} & \longrightarrow & \begin{array}{c} \text{Diagram 6: A square divided into four quadrants by a horizontal and vertical line. A circle is in the bottom-left quadrant, connected to the horizontal line by a curved segment.} \end{array} \quad (2.4)
 \end{array}$$

but not of the ribbon graph they came from. Here additional lines on annulus, pants and torus start from the chosen points on the boundary. The additional lines on the left hand side go to the additional lines in the right hand side under the homeomorphism  $R, \omega$  or  $S$ , which completely determines its isotopy class. Indeed, when the surface (annulus, sphere  $S^2$  with 3 holes, or torus with 1 hole represented by a square with identified edges) is cut along additional lines it becomes a disk, and one knows that a homeomorphism of the boundary of the disk can be extended to a homeomorphism of the disk unique up to isotopy.

$R$  is called inverse Dehn twist, or shortly twist,  $\omega$  is called braiding,  $S$  is called switch. Our first goal will be to give a presentation of the category  $\mathcal{S}$  over the category  $\mathcal{RG}$ , i.e. to add new generators of type  $R, \omega, S$  to  $\mathcal{RG}$  and to find relations, which give a category, equivalent to  $\mathcal{S}$ .

When we present a symmetric monoidal category with left cancellations  $\mathcal{D}$  by generators and relations we mean the following: take a free category  $\mathcal{F}$  on given generators; consider a minimal multiplicative equivalence relation  $\sim$  in  $\text{Mor } \mathcal{F}$  such that in  $\mathcal{F}/\sim$  given relations are satisfied; enlarge the multiplicative equivalence relation  $\sim$  to  $\sim_1$  so

that  $\mathcal{F}/\sim_1$  were symmetric monoidal category for a priori given tensor product; enlarge the multiplicative equivalence relation  $\sim_1$  to  $\sim_2$  so that in  $\mathcal{D} = \mathcal{F}/\sim_2$  any equation  $gf_1 = gf_2$  with  $g, f_1, f_2 \in \text{Mor } \mathcal{D}$  would imply  $f_1 = f_2$ .

**Definition 2.5.** Let  $RG$  be a symmetric monoidal category with left cancellations having ribbon graphs as objects and with morphisms generated over  $\mathcal{RG}$  by new morphisms described as follows.

Let  $\gamma : [0, 1] \hookrightarrow X$  be a curve in a ribbon graph  $(X, B, L)$  such that  $\gamma([0, 1]) \subset \text{Int } X$ ,  $\gamma(0), \gamma(1) \in \partial X - B$ . Such curve will be called a cut. We associate with  $\gamma([0, 1])$  a new morphism  $Tw_\gamma : X \rightarrow X$  called twist.

Let  $h : D_3 \hookrightarrow X$  be a continuous orientation preserving map such that  $h(\partial D_3 - B_3) \subset \partial X - B$ , where hexagon ribbon graph  $(D_3, B_3, L_3)$  is from (2.1) with distinguished interval  $C$ . We associate with  $h$  a new morphism  $Br_h : X \rightarrow X'$  called braiding. Here  $X'$  is obtained from  $X$  by a surgery. Denote by  $A, E$  subintervals of  $B_3$  labeled by  $A, E$ . Cut  $X$  along  $h(A)$  and  $h(E)$  and reglue the edges cross-wise preserving the orientation.

Let  $j : A_1 \hookrightarrow X$  be a continuous orientation preserving map such that  $j(\partial A_1 - B_1) \subset \partial X - B$ , where annulus ribbon graph  $(A_1, B_1, L_1)$  is from (2.2). We associate with it a new morphism  $S_j : X \rightarrow X$  called switch.

These generators are subject to the following relations. If the curve  $\gamma : [0, 1] \hookrightarrow X$  is shrinkable in the class of such curves that  $\gamma(0), \gamma(1) \in \partial X - B$ , we set

$$Tw_\gamma = \text{id}_X.$$


If the map  $h : D_3 \hookrightarrow X$  can be deformed inside the class of maps with condition  $h(\partial D_3 - B_3) \subset \partial X - B$  so that  $h(A)$  or  $h(E)$  shrinks to one point we set

$$Br_h = \phi : X \rightarrow X'$$

if  $h(C)$  shrinks to one point we set

$$Br_h = Tw_\gamma^{-1} \cdot \phi : X \rightarrow X'$$

where  $\phi : X \rightarrow X' \in \mathcal{RG}$  is the unique up to isotopy homeomorphism between  $X$  and  $X'$  which is identity outside of the region  $h(D_3) \cup F$



and  $\gamma : [0, 1] \hookrightarrow h(D_3)$  is a curve isotopic to one of non-shrinkable  $h(A), h(E), h(C)$ .

For each glueing  $g : X \rightarrow Y$  and a map  $\gamma : [0, 1] \hookrightarrow X, h : D_3 \hookrightarrow X$  or  $j : A_1 \hookrightarrow X$  the following diagrams commute:

$$\begin{array}{ccc}
 X & \xrightarrow{Tw_\gamma} & X \\
 g \downarrow & & \downarrow g \\
 Y & \xrightarrow{Tw_{\gamma g}} & Y
 \end{array} \quad (2.5)$$

$$\begin{array}{ccc}
 X & \xrightarrow{Br_h} & X' \\
 g \downarrow & & \downarrow g' \\
 Y & \xrightarrow{Br_{hg}} & Y'
 \end{array} \quad (2.6)$$

$$\begin{array}{ccc}
 X & \xrightarrow{S_j} & X \\
 g \downarrow & & \downarrow g \\
 Y & \xrightarrow{S_{jg}} & Y
 \end{array} \quad (2.7)$$

where the glueing  $g'$  is defined as composition of surgery in  $h(A), h(B)$ , glueing  $g$  and surgery in  $g \circ h(A), g \circ h(B)$ . In these diagrams  $g, g'$  denote isotopy classes of glueings  $g, g'$ . Particular case when  $g$  is isotopic to identity shows that  $Tw_\gamma, Br_h, S_j$  depend only on isotopy classes of  $\gamma, h, j$ .

In the case  $X = D_3$  there are the following relations

$$Br_{BC}^2 = Tw_A Tw_B^{-1} Tw_C^{-1} : \quad \begin{array}{c} \text{Diagram 1} \end{array} \longrightarrow \begin{array}{c} \text{Diagram 2} \end{array} \quad (2.8)$$



$$Tw_A^{-1} = \left( \begin{array}{c} \begin{array}{ccc} \text{Hexagon 1} & \xrightarrow{Br_{AB}} & \text{Hexagon 2} \rightarrow \\ \text{A} & & \text{B} \end{array} \\ \xrightarrow{Br_{AC}} \begin{array}{ccc} \text{Hexagon 3} & \xrightarrow{rot} & \text{Hexagon 4} \\ \text{B} & & \text{A} \end{array} \end{array} \right) \quad (2.9)$$

$$Tw_B^{-1} = \left( \begin{array}{c} \begin{array}{ccc} \text{Hexagon 1} & \xrightarrow{Br_{AB}} & \text{Hexagon 2} \rightarrow \\ \text{A} & & \text{B} \end{array} \\ \xrightarrow{Br_{CB}} \begin{array}{ccc} \text{Hexagon 3} & \xrightarrow{rot^{-1}} & \text{Hexagon 4} \\ \text{C} & & \text{A} \end{array} \end{array} \right) \quad (2.10)$$

Here  $Br_{AB}$  stands for  $Br_{\text{id}}$ , similarly  $Br_{AC}, Br_{CD}$  are defined and  $rot$  denotes a morphism from  $\mathcal{RG}$ , rotation by  $2\pi/3$ .

There are two relations for  $X = D_4 =$  a disk with 4 intervals marked on the boundary:

$$\begin{array}{ccc} \begin{array}{c} \text{Disk 1} \\ \text{Z} \text{ (top-left), } Y \text{ (top-right)} \\ \text{A} \text{ (bottom-left), } B \text{ (bottom-right)} \\ \text{z}Br_{BC}^{\pm 1} \searrow \end{array} & \xrightarrow{D Br_{YC}^{\pm 1}} & \begin{array}{c} \text{Disk 2} \\ \text{Y} \text{ (top-left), } U \text{ (top-right)} \\ \text{C} \text{ (bottom-left), } A \text{ (bottom-right)} \\ \nearrow U Br_{AC}^{\pm 1} \end{array} \\ & & \begin{array}{c} \text{Disk 3} \\ \text{Z} \text{ (top-left), } U \text{ (top-right)} \\ \text{A} \text{ (bottom-left), } B \text{ (bottom-right)} \end{array} \end{array} \quad (2.11)$$

Here three inclusions  $D_3 \hookrightarrow D_4$  are determined by 3 indices appended to braidings. Image of  $D_3$  covers a half of  $D_4$ .

Relations for  $X = A_1$  are

$$(ST)^3 = S^2 : \quad \begin{array}{c} \text{Diagram 1} \end{array} \longrightarrow \begin{array}{c} \text{Diagram 2} \end{array}, \quad (2.12)$$

where  $T = Tw_\gamma$ , and (2.13) (see Figure 1).

$$\begin{array}{ccc} \begin{array}{c} X \\ \text{Diagram 1} \end{array} & \xrightarrow{S^2} & \begin{array}{c} X \\ \text{Diagram 2} \end{array} \\ \downarrow \text{\scriptsize } {}_X Br_{\gamma\gamma}^{-1} & & \parallel \\ \begin{array}{c} X \\ \text{Diagram 3} \end{array} & \xrightarrow{Tw_\gamma^{-1}} & \begin{array}{c} X \\ \text{Diagram 4} \end{array} \end{array} \quad (2.13)$$

FIGURE 1. A relation for annulus ribbon graph  $A_1$

Finally, there is a relation for annulus ribbon graph  $A_2$  with two intervals marked on the boundary (see diagram 2.14, Figure 2). Here  $rot_\pi \in \mathcal{RG}$  is the rotation by  $\pi$ , and  $\phi \in \mathcal{RG}$  is a homeomorphism, which is identity in a neighbourhood of the hole and slides intervals  $X, Y$  along the boundary.

*Remark 2.1.* The requirement that  $RG$  is a symmetric monoidal category with left cancellations produces more relations than just in a category with relations (2.5)–(2.14). For instance, any two generators  $f_1, f_2$  with essentially non-intersecting supports  $(\gamma([0, 1]), h(D_3), j(A_1))$  commute (this can be called a “locality principle”). Essentially non-intersecting means that images of some  $\gamma', h', j'$  isotopic to  $\gamma, h, j$  do


FIGURE 2. A relation for annulus ribbon graph  $A_2$ 

not intersect. Indeed, cut the ribbon graph  $X$  into pieces so that one of the pieces contained support of  $f_1$  and another contained support of  $f_2$ . Denote the resulting ribbon graph  $\tilde{X}$ . The liftings  $\tilde{f}_1, \tilde{f}_2 : \tilde{X} \rightarrow \tilde{X}$  commute because they are tensor products (disjoint unions) of a generator with identity map. The axioms (2.5)–(2.7) imply that  $gf_1f_2 = gf_2f_1 : \tilde{X} \rightarrow \tilde{X}$ . The left cancellation property implies that  $f_1, f_2 : X \rightarrow X$  commute.


Also (2.8)–(2.14) imply relations of the type (2.8)–(2.14) in  $X$  associated with embeddings of  $D_3, D_4, A_1, A_2$  into  $X$ .

**Proposition 2.2.** *The duplication functor  $\text{dupl} : \mathcal{RG} \rightarrow \mathcal{S}$  extends to a functor  $\text{dupl} : RG \rightarrow \mathcal{S}$ . It sends  $Tw_{\gamma} : X \rightarrow X$  to the inverse*

Dehn twist  $R$  in  $X \cup \bar{X}$  performed in neighbourhood of the cycle  $\gamma \cup \bar{\gamma}$ . The braiding  $Br_h : X \rightarrow X'$  goes to a braiding homeomorphism  $\omega : X \cup \bar{X} \rightarrow X' \cup \bar{X}'$ , supported in  $h(D_3) \cup h(\bar{D}_3)$ , looking as (2.3)



(exterior of the first figure is  $h(\bar{D}_3)$ ). The switch  $S_j : X \rightarrow X$  goes to a switching homeomorphism  $S : X \cup \bar{X} \rightarrow X \cup \bar{X}$  supported in  $j(A_1) \cup j(\bar{A}_1)$  looking as (2.4)



(the upper half of the square with identified edges is  $j(A_1)$ , the lower half is  $j(\bar{A}_1)$ ).

### 3. Ribbon graphs compared to surfaces

We say that a ribbon graph  $(X, B)$  has  $g$  loops with  $n$  entries if  $\dim H_1(X, \mathbb{R}) = g$ ,  $\text{Card } \pi_0(B) = n$ . Duplicated, it gives a surface  $X \cup \bar{X}$  of genus  $g$  with  $n$  holes.

**Theorem 3.1.** *There exists a functor  $\Phi : \text{Surf} \rightarrow \text{RG}$  such that the composition  $\text{dupl} \circ \Phi : \text{Surf} \rightarrow \mathcal{S}$  is isomorphic to the inclusion functor  $\text{Surf} \hookrightarrow \mathcal{S}$ .*

*Proof.* The theorem reduces to constructing a splitting of the homomorphism

$$\text{Aut}_{\text{RG}}(Y_{g,n}) \rightarrow \text{Aut}_{\mathcal{S}}(\text{dupl}(Y_{g,n})) \simeq \text{Aut}_{\text{Surf}}(\text{dupl}(Y_{g,n})).$$

Here  $Y_{g,n}$  means a graph with  $g$  loops (of genus  $g$ ) and with  $n$  components of  $B$  (entries). In general,  $Y_{g,n}$  is labeled by  $n_1$  copies of  $L_1 \in \mathcal{C}$ ,  $n_2$  copies of  $L_2 \in \mathcal{C}$ , etc. Each morphism induces a permutation of the set  $\pi_0(B) \simeq \pi_0(\partial(\text{dupl}(Y_{g,n})))$ , so there are group homomorphisms


$$\text{Aut}_{\text{RG}}(Y_{g,n}) \rightarrow \text{Aut}_{\mathcal{S}}(\text{dupl}(Y_{g,n})) \rightarrow \mathfrak{S}_n.$$

In particular case when all labels coincide, the automorphism groups  $G_{RG}, G_S$  will be the biggest, and the general case groups are just preimages under projections  $G_{RG} \rightarrow \mathfrak{S}_n, G_S \rightarrow \mathfrak{S}_n$  of the subgroup of permutations leaving invariant  $n_k$ -element subset of  $\pi_0(B)$  with label  $L_k, k \geq 1, n = n_1 + n_2 + \dots$ . Thus it suffices to consider the case when all labels  $L \in \mathcal{C}$  coincide. We have to construct a splitting of the homomorphism

$$\text{Aut}_{RG}(Y_{g,n}) \rightarrow \text{Aut}_S(\text{dupl}(Y_{g,n})) = M_{g,n},$$

where the last group is the mapping class group of a surface of genus  $g$  with  $n$  holes.

Using braiding isomorphisms any ribbon graph  $Y_{g,n}$  can be reduced to a standard form



so we have to consider only standard case  $Y_{g,n} = X_{g,n}$ . Using results of Dehn [2] and Lickorish [7, 8], Birman [1] gave a set of generators of  $M_{g,n}$ . These consist of braidings permuting the holes, some Dehn twists coming from  $Tw_\gamma \in RG$ , and some Dehn twists along cycles in  $\text{dupl}(X_{g,n})$  which can be obtained from cycles of the form  $\gamma \cup \bar{\gamma}$  by action of homeomorphisms  $S$ . This implies that the map

$$\text{Aut}_{RG}(X_{g,n}) \rightarrow M_{g,n}$$

is an epimorphism. More details will be given below.

When  $g \leq 1$  or  $n \leq 1$  not only generators of  $M_{g,n}$  but also relations are known (Magnus [14] for  $g = 0$ , Birman [1] for  $g = 1$  and Wajnryb [22] for  $n = 0, 1$ ). For generic values of  $g, n$  defining relations were not written, but it is known (Birman [1], Scott [19]) that there is an exact sequence

$$1 \rightarrow \bar{B}_n(\Sigma_g) \rightarrow M_{g,n} \rightarrow M_{g,0} \rightarrow 1,$$

where  $\bar{B}_n(\Sigma_g)$  is the extension of the braid group of a closed surface of genus  $g$  by  $\mathbb{Z}^n$ . Its defining relations are found by Scott [19]. This information is sufficient to show case by case that generators of  $M_{g,n}$  admit liftings to  $\text{Aut}_{RG}(X_{g,n})$  and the relations in  $M_{g,n}$  lift to relations in  $\text{Aut}_{RG}(X_{g,n})$ . This implies the theorem.  $\square$

The topological version of the Grothendieck conjecture about Teichmüller's tower is

*Conjecture .* The duplication functor  $dupl : RG \rightarrow \mathcal{S}$  is an equivalence of categories.

## 4. A category of trivalent nets

### 4.1. Trivalent nets.

**Definition 4.1.** Let a trivalent net  $\Gamma$  be a 1-complex with set of vertices  $V \sqcup B$  and set of edges  $E$ . Elements of  $V$ , called 3-vertices, occur thrice as endpoints of edges, and elements of  $B$ , called ends, occur once as an endpoint of an edge. Each edge has at least one endpoint in  $V$  and  $\Gamma$  is equipped with labeling of ends  $B \rightarrow \mathcal{C}$ . For each vertex  $v \in V$  belonging to 3 edges  $a, b, c$  denote  $a', b', c'$  3 different isotopy classes of embeddings  $\gamma : [0, 1] \hookrightarrow a, b, c$  such that  $\gamma(0) = v$ . A cyclic order in the set  $\{a', b', c'\}$  is chosen (which is one of the two cyclic permutations  $(a', b', c')$  or  $(a', c', b')$ ).


An edge connecting a 3-vertex and an end is called external leg, others are internal edges. Trivalent nets will be drawn in the plane in such a way that cyclic orientation of edges around a 3-vertex is coherent with the orientation of the plane of drawing.

Let  $\Gamma_1, \Gamma_2$  be trivalent nets.

**Definition 4.2.** A glueing  $g : \Gamma_1 \rightarrow \Gamma_2$  is a surjective continuous mapping of 1-complexes (sending vertices to vertices and edges into edges), bijective on 3-vertices and preserving cyclic order for edges incident to each 3-vertex, and such that preimage of an end is an end with the same label from  $\mathcal{C}$ , preimage of an internal edge is an internal edge or a pair of external legs with dual labels.

Denote  $\mathcal{TN}_0$  the category of trivalent nets with glueings as morphisms. It is a symmetric monoidal category with disjoint union as a monoidal product.

**Definition 4.3.** Let  $\mathcal{TN}$  be a symmetric monoidal category with left cancellations having trivalent nets as objects and morphisms, generated over  $\text{Mor } \mathcal{TN}_0$  by fusing moves: for each internal edge  $a \in \Gamma \in \text{Ob } \mathcal{TN}_0$  with different endpoints the fusing move  $fus_a : \Gamma \rightarrow \Gamma'$  is a new morphism, where  $\Gamma'$  differs from  $\Gamma$  in a neighbourhood of  $a$  as shown in the figure



New morphisms are subject to the following relations (commutative diagrams). For any glueing  $g$  and any internal edge  $a \in \Gamma_1$

$$\begin{array}{ccc} \Gamma_1 & \xrightarrow{fus_a} & \Gamma'_1 \\ g \downarrow & & \downarrow g' \\ \Gamma_2 & \xrightarrow{fus_{g(a)}} & \Gamma'_2 \end{array} \quad (4.1)$$

The following relations are written literally, without any complementary part


$$\left( \begin{array}{c} \text{Diagram 1} \xrightarrow{fus_a} \text{Diagram 2} \xrightarrow{fus_b} \text{Diagram 3} \end{array} \right) = \text{id} \quad (4.2)$$

and (4.3), Figure 3.

(4.3)

FIGURE 3. The pentagon relation

There is a functor  $fat : \mathcal{TN} \rightarrow \mathcal{RG}$  called fattening. Given a trivalent net  $\Gamma$  with a set of 3-vertices  $V$ , we take  $\text{Card } V$  copies of a ribbon graph  $D_3$ , glue pairwise intervals corresponding to internal edges of  $\Gamma$ , and put on  $fat(\Gamma)$  the same labels as on  $\Gamma$ . It is important to glue respecting the orientation of  $D_3$ 's, and to make a bijection between  $\pi_0(B_3)$  and the set of edges incident to a 3-vertex respecting the cyclic ordering ( $\pi_0(B_3)$  has a canonical cyclic ordering coming from the orientation of  $D_3$ ). Fattening of a glueing is a glueing and fattening of the fusing move  $fus_a$  is the isotopy class of a homeomorphism  $fat(\Gamma) \rightarrow fat(\Gamma')$ , which is identity outside of the two  $D_3$ 's glued by  $a$ .



The ribbon graphs  $fat(\Gamma)$  are canonically marked. A *marking*  $M$  of a ribbon graph  $(X, B)$  is by definition an isotopy class of  $\bigcup_k \gamma_k([0, 1])$  for non-intersecting cuts  $\gamma_k : [0, 1] \hookrightarrow X$  (such that  $\gamma_k([0, 1]) \subset \text{Int } X$ ,  $\gamma_k(0), \gamma_k(1) \in \partial X - B$ ) and such that  $X - \bigcup_k \gamma_k([0, 1])$  is a union of hexagons  $D_3$ .

#### 4.2. Coherence theorem for markings of ribbon graphs.

**Theorem 4.1.** *Any marking of a ribbon graph can be obtained from any other marking by a sequence of fusing moves. Any such sequence can be transformed into any other using the pentagon, quadrilateral and  $fus^2 = \text{id}$  identity.*

**Corollary 4.2.** *For all trivalent nets  $\Gamma_1, \Gamma_2$*

$$\mathcal{TN}(\Gamma_1, \Gamma_2) \rightarrow \mathcal{RG}(fat(\Gamma_1), fat(\Gamma_2))$$

*is bijective.*

*Proof.* Any morphism  $f$  in  $\mathcal{TN}$  or  $\mathcal{RG}$  is uniquely decomposed as  $gh$ , where  $g$  is a canonical glueing determined by the set of pairs of ends glued by  $f$ , and  $h$  is an isomorphism. Hence, we can consider only isomorphisms. Let  $\phi : fat(\Gamma_1) \rightarrow fat(\Gamma_2)$  be a homeomorphism. Use it to transfer the canonical marking of  $fat(\Gamma_2)$  to a marking  $M$  of  $fat(\Gamma_1)$ . By Theorem it can be obtained from the canonical marking of  $fat(\Gamma_1)$  by a sequence of fusing moves. This sequence determines a morphism  $j : \Gamma_1 \rightarrow \Gamma \in \mathcal{TN}$  where  $\Gamma$  is the net determined by the marking  $M$ .



It is identified with  $\Gamma_2$  by an isomorphism  $i : \Gamma \rightarrow \Gamma_2 \in \mathcal{TN}_0$ . Finally, for  $h = ji$  we have  $\phi = fat(h)$ .

For the second statement, it suffices to consider the case of isomorphic  $\Gamma_1$  and  $\Gamma_2$ . Furthermore, we have only to prove that any  $f : \Gamma \rightarrow \Gamma \in \mathcal{TN}$ , such that  $fat(f)$  is isotopic to  $id_{fat(\Gamma)}$ , equals  $id_\Gamma$ . Represent  $f$  as a product  $fus_{a_1} fus_{a_2} \dots fus_{a_n}$ . The sequence of fusing moves in  $fat(\Gamma)$  corresponds to it. This sequence produces a new marking of  $fat(\Gamma)$ , which equals to the image of the canonical marking under the homeomorphism  $fat(f)^{-1}$ . Hence, it is isotopic to the canonical marking. This gives a loop in  $\mathcal{M}$  which by the Theorem can be shrunk by the pentagon, quadrilateral and  $fus^2 = id$  moves. Parallely we reduce the expression for  $f$  to 1, using the pentagon, quadrilateral and  $fus^2 = id$  identities in  $\mathcal{TN}$ .  $\square$

This corollary implies

**Theorem 4.3.** *The functor  $fat : \mathcal{TN} \rightarrow \mathcal{RG}$  is an equivalence of  $\mathcal{TN}$  with a full subcategory  $\mathcal{RG}_+$  of ribbon graphs, each component of which satisfy the inequalities*

$$n \geq 3 \quad \text{or} \quad g \geq 1, n \geq 1 \quad \text{or} \quad g \geq 2 \quad (4.4)$$

( $g, n$  stand for genus and number of marked intervals).

*Remark 4.1.* Formally adjoining the objects  $D_0, D_1, D_2, A_0$  and some morphisms to  $\mathcal{RG}_+$  we can reconstruct  $\mathcal{RG}$ . We shall do a similar thing later.

**4.3. Trivalent nets with twists, braiding and switching.** Now we add to  $\mathcal{TN}$  new morphisms just as we added them to  $\mathcal{RG}$ . The obtained category  $TN$  will be a full subcategory of  $RG$ .

**Definition 4.4.** Let  $TN$  be a symmetric monoidal category with left cancellations which objects are trivalent nets and morphisms are generated over  $\mathcal{TN}$  by the following morphisms.

For an edge  $A$  of a trivalent net  $\Gamma$  there is a twist automorphism  $Tw_A : \Gamma \rightarrow \Gamma$ .

For a 3-vertex  $a \in \Gamma$  and two incident edges  $B, C$  there is a braiding morphism  ${}_a Br_{BC} : \Gamma \rightarrow \Gamma'$ , denoted also  ${}_A Br_{BC}$  or  $Br_{BC}$  if there is no ambiguity, where  $\Gamma'$  is obtained from  $\Gamma$  by a surgery cutting edges  $B$  and  $C$  and reglueing the ends crosswise.  $\Gamma'$  is isomorphic to  $\Gamma$  with the changed cyclic orientation at  $a$ . Locally it looks like

$${}_a Br_{BC} : \begin{array}{c} A \\ | \\ \bullet \\ / \quad \backslash \\ B \quad C \end{array} \longrightarrow \begin{array}{c} A \\ | \\ \bullet \\ \backslash \quad / \\ C \quad B \end{array}$$

For each loop-edge  $A$  of  $\Gamma$  there is a switch automorphism  $S_A : \Gamma \rightarrow \Gamma$ , e.g.

$$S_A : \text{---} \bullet \bigcirc^A \rightarrow \text{---} \bullet \bigcirc^A.$$

The morphisms are subject to the following relations. For any glueing  $g$

$$\begin{array}{ccc} \Gamma_1 & \xrightarrow{Tw_A} & \Gamma_1 \\ g \downarrow & & \downarrow g \\ \Gamma_2 & \xrightarrow{Tw_{g(A)}} & \Gamma_2 \end{array} \quad (4.5)$$

$$\begin{array}{ccc} \Gamma_1 & \xrightarrow{aBr_{BC}} & \Gamma'_1 \\ g \downarrow & & \downarrow g' \\ \Gamma_2 & \xrightarrow{g(a)Br_{g(B),g(C)}} & \Gamma'_2 \end{array} \quad (4.6)$$

$$\begin{array}{ccc} \Gamma_1 & \xrightarrow{S_A} & \Gamma_1 \\ g \downarrow & & \downarrow g \\ \Gamma_2 & \xrightarrow{S_{g(A)}} & \Gamma_2 \end{array} \quad (4.7)$$

commute. Also

$$Br_{BC}^2 = Tw_A Tw_B^{-1} Tw_C^{-1} : \begin{array}{c} A \\ | \\ \bullet \\ / \quad \backslash \\ B \quad C \end{array} \longrightarrow \begin{array}{c} A \\ | \\ \bullet \\ / \quad \backslash \\ B \quad C \end{array} \quad (4.8)$$

$$Tw_A^{-1} = \left( \begin{array}{c} C \\ | \\ \bullet \\ / \quad \backslash \\ A \quad B \end{array} \xrightarrow{Br_{AB}} \begin{array}{c} C \\ | \\ \bullet \\ / \quad \backslash \\ B \quad A \end{array} \xrightarrow{Br_{AC}} \begin{array}{c} A \\ | \\ \bullet \\ / \quad \backslash \\ B \quad C \end{array} \xrightarrow{rot} \begin{array}{c} C \\ | \\ \bullet \\ / \quad \backslash \\ A \quad B \end{array} \right) \quad (4.9)$$

$$Tw_B^{-1} = \left( \begin{array}{c} C \\ | \\ \bullet \\ / \quad \backslash \\ A \quad B \end{array} \xrightarrow{Br_{AB}} \begin{array}{c} C \\ | \\ \bullet \\ / \quad \backslash \\ B \quad A \end{array} \xrightarrow{Br_{CB}} \begin{array}{c} B \\ | \\ \bullet \\ / \quad \backslash \\ C \quad A \end{array} \xrightarrow{rot^{-1}} \begin{array}{c} C \\ | \\ \bullet \\ / \quad \backslash \\ A \quad B \end{array} \right) \quad (4.10)$$

(here  $rot$  is an isomorphism of rotation),

$$\begin{array}{ccccc}
 \begin{array}{c} X \quad C \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ A \quad B \end{array} & \xrightarrow{Br_{BC}^{\pm 1}} & \begin{array}{c} X \quad B \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ A \quad C \end{array} & \xrightarrow{fus} & \begin{array}{c} X \quad B \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ A \quad C \end{array} \\
 \downarrow fus & & & & \downarrow Br_{AC}^{\pm 1} \\
 \begin{array}{c} X \quad C \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ A \quad B \end{array} & \xrightarrow{Br_{YC}^{\pm 1}} & \begin{array}{c} X \quad B \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ C \quad A \end{array} & \xrightarrow{fus} & \begin{array}{c} X \quad B \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ C \quad A \end{array} \\
 & & & & (4.11)
 \end{array}$$

$$(S_M Tw_M)^3 = S_M^2 : \text{---} \bigcirc^M \text{---} \rightarrow \text{---} \bigcirc^M \text{---}, \quad (4.12)$$

$$\begin{array}{ccc}
 \begin{array}{c} X \quad M \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \end{array} & \xrightarrow{S_M^2} & \begin{array}{c} X \quad M \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \end{array} \\
 \downarrow {}_X Br_{MM}^{-1} & & \parallel \\
 \begin{array}{c} X \quad M \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \end{array} & \xrightarrow{Tw_M^{-1}} & \begin{array}{c} X \quad M \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \end{array} \\
 & & (4.13)
 \end{array}$$

and (4.14) (see Figure 4) are satisfied.

It is clear that the functor  $fat : \mathcal{TN} \rightarrow \mathcal{RG}$  extends to a functor  $fat : TN \rightarrow RG$ .


**Theorem 4.4.** *The functor  $fat : TN \rightarrow RG$  is an equivalence of  $TN$  with a full subcategory  $RG_+$  of ribbon graphs satisfying condition (4.4).*

*Proof.* A quasiinverse functor  $\Phi : \mathcal{RG}_+ \rightarrow \mathcal{TN}$  is constructed like that. For any ribbon graph  $X$  choose a marking  $M_X$  and set  $\Phi(X)$  to be the trivalent net  $N(M_X)$  associated with  $M_X$ . Let  $g : X \rightarrow Y$  be a glueing, then there is a marking  $g^*(M_X)$  of  $Y$  consisting of  $g(M_X)$  and images of those boundary intervals of  $X$  which are glued by  $g$ . To  $g$  corresponds the morphism

$$\Phi(X) = N(M_X) \xrightarrow{g'} N(g^*(M_X)) \xrightarrow{f} N(M_Y) = \Phi(Y),$$

where  $g'$  is a canonical glueing, and  $f$  is a composition of fusings, transforming  $g^*(M_X)$  into  $M_Y$ .

The functor  $\Phi$  extends to a functor  $\Phi : RG_+ \rightarrow TN$ . Indeed, take a map  $\gamma : [0, 1] \hookrightarrow X$ , or  $h : (D_3, B_3) \hookrightarrow X$ , or  $j : (A_1, B_1) \hookrightarrow X$  which

FIGURE 4. A relation for  $A_2$  net

determines a morphism  $Tw_\gamma$ ,  $Br_h$ , or  $S_j$  as in Definition 2.5. There exists a marking  $M$  of  $X$  containing  $\gamma([0, 1])$ , or  $h(B_3)$ , or  $j(B_1)$  and a composition  $f$  of fusing moves relating  $M_X$  and  $M$ . Set  $\Phi(Tw_\gamma) = f Tw_A f^{-1}$ , or  $\Phi(S_j) = f S_A f^{-1}$  for  $A$  from  $M$  determined by  $\gamma$  or  $h$  and similarly for  $Br_h$ . The morphism  $f \in \mathcal{TN}$  is unique and a different choice of marking  $M$ , say  $M'$ , will give  $f' = fg$ , where  $g$  is a composition of fusing moves  $fus_a$  with  $a \neq \gamma([0, 1])$ ,  $a \notin h(B_3)$ ,  $a \neq j(B_1)$ . Hence,  $\Phi(Tw_\gamma)$ ,  $\Phi(Br_h)$ ,  $\Phi(S_j)$  are correctly defined morphisms of  $TN$ .

The relations (4.5)–(4.7) follow from the relations (2.5)–(2.7), and  $\Phi$  applied to (4.8)–(4.14) is just (2.8)–(2.14). Thus the functor  $\Phi$  :

$RG_+ \rightarrow TN$  is constructed and one can easily see that it is quasiinverse to  $fat : TN \rightarrow RG_+$ .  $\square$

## 5. A category of nets

5.1. **Nets.** Now we consider nets with valency not bigger than 3.

**Definition 5.1.** Let a net be a 1-complex  $\Gamma$  with the set of vertices  $V_3 \sqcup V_2 \sqcup V_1 \sqcup B$  and the set of edges  $E$ . The elements of  $V_i$ , called  $i$ -vertices ( $i = 1, 2, 3$ ) occur  $i$  times as endpoints of edges, elements of  $B$  called ends occur once as an endpoint of an edge. We assume that each edge has at least one endpoint in  $V_3 \sqcup V_2 \sqcup V_1$ . A net  $\Gamma$  is equipped with a labeling of ends  $B \rightarrow \mathcal{C}$  and for each 3-vertex a cyclic order in the set of incident edges is chosen.

Denote  $\mathcal{N}_0$  the symmetric monoidal category of nets with glueings as morphisms. Glueings are defined similarly to trivalent case with requirement that a glueing is bijective on  $i$ -vertices,  $i = 1, 2, 3$ .

**Definition 5.2.** Let  $\mathcal{N}$  be a symmetric monoidal category with left cancellations which has nets as objects, and morphisms are generated over  $\mathcal{N}_0$  by the following morphisms:

insertion  $ins_a : \Gamma \rightarrow \Gamma'$  is an isomorphism of  $\Gamma$  to  $\Gamma'$  with a 1-vertex  $a \in \Gamma'$  and the incident edge deleted or a 2-vertex  $a \in \Gamma'$  deleted and the two incident edges combined in one;

deletion  $del_a : \Gamma' \rightarrow \Gamma$  is the inverse morphism.

These morphisms are subject to the following relations. Whenever the diagram

$$\begin{array}{ccc} \Gamma_1 & \xrightarrow{ins_a} & \Gamma'_1 \\ g \downarrow & & \downarrow g' \\ \Gamma_2 & \xrightarrow{ins_{g'(a)}} & \Gamma'_2 \end{array} \quad (5.1)$$

commutes as a diagram of mappings of 1-complexes, it commutes also in  $\mathcal{N}$ . Also the diagrams

$$\begin{array}{ccc} \begin{array}{c} \text{---} a \bullet \end{array} & \xrightarrow{ins_b} & \begin{array}{c} \text{---} a \bullet \text{---} b \bullet \end{array} \\ ins_c \downarrow & & \downarrow ins_c \\ \begin{array}{c} \text{---} a \bullet \text{---} c \bullet \end{array} & \xrightarrow{ins_b} & \begin{array}{c} \text{---} a \bullet \begin{array}{l} \nearrow c \bullet \\ \searrow b \bullet \end{array} \end{array} \end{array} \quad (5.2)$$

$$\begin{array}{ccc}
\begin{array}{c} \bullet \\ \hline a \end{array} & \xrightarrow{ins_b} & \begin{array}{c} \bullet \quad \bullet \\ \hline b \quad a \end{array} \\
ins_c \downarrow & \nearrow \phi & \\
\begin{array}{c} \bullet \quad \bullet \\ \hline a \quad c \end{array} & & 
\end{array} \tag{5.3}$$

where  $\phi$  is the isomorphism of nets,

$$\begin{array}{ccc}
\begin{array}{c} \bullet \\ \hline a \end{array} & \xrightarrow{ins_b} & \begin{array}{c} \bullet \quad \bullet \\ \hline b \quad a \end{array} \\
ins_c \downarrow & \nearrow \psi & \\
\begin{array}{c} \bullet \quad \bullet \\ \hline a \quad c \end{array} & & 
\end{array} \tag{5.4}$$

where  $\psi$  is the isomorphism of nets sending  $a \mapsto b$ ,  $c \mapsto a$ , are all commutative.

Finally, the isomorphism

$$rot : \begin{array}{c} \bullet \quad \bullet \\ \hline \end{array} \longrightarrow \begin{array}{c} \bullet \\ \hline \bullet \end{array} \tag{5.5}$$

interchanging the vertices is set equal to identity morphism.

Let  $\mathcal{N}_+$  be the full subcategory of such nets that genus  $g$  and the number of ends  $n$  for each component satisfy (4.4). There is an obvious inclusion functor  $inc : \mathcal{TN}_0 \rightarrow \mathcal{N}_+$ . Let us construct a functor  $tri : \mathcal{N}_+ \rightarrow \mathcal{TN}_0$ . Take a net  $\Gamma$ . Denote  $red_1(\Gamma)$  the net  $\Gamma$  with all 1-vertices and incident edges deleted. Repeat the reduction until  $\Gamma' = (red_1)^k(\Gamma)$  does not contain 1-vertices. Change each connected string of 2-vertices in  $\Gamma'$  together with incident edges to one edge. The obtained net  $tri(\Gamma)$  is trivalent. Any glueing  $g : \Gamma_1 \rightarrow \Gamma_2$  is bijective on 1-vertices, hence, induces a glueing  $red_1(\Gamma_1) \rightarrow red_1(\Gamma_2)$ , therefore, a glueing  $\Gamma'_1 \rightarrow \Gamma'_2$  which leads to a glueing  $tri(g) : tri(\Gamma_1) \rightarrow tri(\Gamma_2)$ . Any morphism  $ins_a$ ,  $del_a$  is sent to identity by the functor  $tri$ .

**Theorem 5.1.** *The functors  $inc : \mathcal{TN}_0 \rightarrow \mathcal{N}_+$  and  $tri : \mathcal{N}_+ \rightarrow \mathcal{TN}_0$  are quasi-inverse to each other.*

**5.2. Nets with fusing, braiding, twists and switches.** Now we add more morphisms to the category of nets.

**Definition 5.3.** Let  $\mathcal{N}$  be a symmetric monoidal category with left cancellations having nets as objects and having morphisms generated over  $\mathcal{N}$  by  $fus_a : \Gamma \rightarrow \Gamma'$ ,  $Tw_a : \Gamma \rightarrow \Gamma$ ,  ${}_a Br_{BC} : \Gamma \rightarrow \Gamma'$ ,  $S_a : \Gamma \rightarrow \Gamma$  as in Definitions 4.3, 4.4. We subject them to the relations of Definitions 4.3, 4.4 and the following. There are equations

$$Tw_A = Tw_B : \begin{array}{c} A \quad B \\ \hline \bullet \end{array} \longrightarrow \begin{array}{c} A \quad B \\ \hline \bullet \end{array}, \tag{5.6}$$

(5.7)

The following diagrams commute

(5.8)

$$(5.9)$$

Denote  $N_+$  a full subcategory of nets whose components satisfy inequalities (4.4).

**Theorem 5.2.** *The functor  $\text{tri}$  uniquely extends to a functor  $\text{tri} : N_+ \rightarrow TN$  quasi-inverse to inclusion functor  $\text{inc} : TN \rightarrow N_+$ .*

**5.3. Nets compared with ribbon graphs.** Extend the functor  $N_+ \xrightarrow{tri} TN \xrightarrow{fat} RG$  to a functor  $fat : N \rightarrow RG$  setting

$$\begin{aligned}
 fat(\bullet \text{---} \bullet) &= \bigcirc \\
 fat(\bullet \text{---} \overline{A}) &= \overline{\bigcirc_A}, \\
 fat(\overline{A} \text{---} \bullet) &= \overline{\bigcirc_B}, \\
 fat(\bullet \bigcirc \bullet) &= \text{a square with rounded corners and a central circle},
 \end{aligned}$$

The fattening of an insertion or deletion morphism is set to the unique “identity” homeomorphism of the ribbon graphs. Fattenings of other generators are already defined.

**Theorem 5.3.** *The functor  $fat : N \rightarrow RG$  is an equivalence of categories.*

## 6. A category of oriented nets

**6.1. Oriented nets.** By an *oriented net* we mean a net  $\Gamma$ , equipped with such orientation of edges that each 2-vertex or 3-vertex is a source for some edge and a target for some edge. Given an oriented net  $\Gamma$  with labeling  $L : B \rightarrow \mathcal{C}$ , we define its desorientation  $deso(\Gamma)$  as a net  $\Gamma$  with labeling  $L' : B \rightarrow \mathcal{C}$  constructed as follows.  $L'(b) = L(b)$  if the end  $b$  is a target and  $L'(b) = L(b)^\sim$  if the end  $b$  is a source of the incident external leg. A glueing of oriented nets is an orientation preserving map, which is a glueing of underlying desoriented nets. In particular, only those external legs can be glued which carry the same label from  $\mathcal{C}$  and one of them has the incident end as a target, and another as a source.

**Definition 6.1.** Let  $\mathcal{ON}$  be a symmetric monoidal category with left cancellations having oriented nets as objects and morphisms generated over glueings by the following new morphisms:

reversal  $rev_a : \Gamma \rightarrow \Gamma'$ , where  $a$  is an internal edge of  $\Gamma$  and  $\Gamma'$  differs from  $\Gamma$  by a reversed orientation of  $a$ ;

dualising morphism  $du_a : \Gamma \rightarrow \Gamma'$ , where  $a$  is an external leg of  $\Gamma$ ,  $\Gamma'$  differs from  $\Gamma$  by a reversed orientation of  $a$ , and the labels  $A, A' \in \mathcal{C}$  attached to  $a$  are dual to each other;

$ins_a : \Gamma \rightarrow \Gamma'$ ,  $del_a : \Gamma \rightarrow \Gamma'$ , insertion and deletion of a 2-vertex.

These morphisms are subject to relations

$$ins \cdot del = 1, \quad del \cdot ins = 1, \quad rev_a^2 = 1, \quad du_a^2 = 1.$$

For each glueing  $g : \Gamma_1 \rightarrow \Gamma_2$  the diagram (5.1) and

$$\begin{array}{ccc} \Gamma_1 & \xrightarrow{rev_a} & \Gamma'_1 \\ g \downarrow & & \downarrow g' \\ \Gamma_2 & \xrightarrow{rev_{g(a)}} & \Gamma'_2 \end{array} \quad (6.1)$$

$$\begin{array}{ccc} \Gamma_1 & \xrightarrow{du_a} & \Gamma'_1 \\ g \downarrow & & \downarrow g' \\ \Gamma_2 & \xrightarrow{du_{g(a)}} & \Gamma'_2 \end{array} \quad (6.2)$$



(if the external leg  $a$  is not glued by  $g$ ) commute. The diagrams

$$\begin{array}{ccc}
 \begin{array}{c} \xrightarrow{\quad} \bullet \xrightarrow{\quad} a \\ \text{\scriptsize $ins_c$} \downarrow \\ \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} c \end{array} & \begin{array}{c} \xrightarrow{\text{\scriptsize $ins_b$}} \\ \nearrow \psi \end{array} & \begin{array}{c} \xrightarrow{\quad} \bullet \xrightarrow{\quad} b \xrightarrow{\quad} \bullet \xrightarrow{\quad} a \\ \end{array}
 \end{array} \quad (6.3)$$

$$\begin{array}{ccc}
 \begin{array}{c} A \diagdown \quad \xrightarrow{X} \quad \xrightarrow{X} \quad \diagup D \\ B \diagup \quad \quad \quad \diagdown C \end{array} & \xrightarrow{\text{\scriptsize $du_X \ du_X$}} & \begin{array}{c} A \diagdown \quad \xrightarrow{X^\sim} \quad \xrightarrow{X^\sim} \quad \diagup D \\ B \diagup \quad \quad \quad \diagdown C \end{array} \\
 \downarrow g' & & \downarrow g'' \\
 \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ B \quad C \end{array} & \xrightarrow{\text{\scriptsize $rev$}} & \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ B \quad C \end{array}
 \end{array} \quad (6.4)$$

$$\begin{array}{ccc}
 \begin{array}{c} \xrightarrow{X} \quad \xrightarrow{X} \quad \diagup B \\ \bullet \quad \quad \quad \diagdown A \end{array} & \xrightarrow{\text{\scriptsize $du_X \ du_X$}} & \begin{array}{c} \xrightarrow{X^\sim} \quad \xrightarrow{X^\sim} \quad \diagup B \\ \bullet \quad \quad \quad \diagdown A \end{array} \\
 \downarrow g' & & \downarrow g'' \\
 \begin{array}{c} \quad \quad \diagup B \\ \bullet \quad \quad \diagdown A \end{array} & \xrightarrow{\text{\scriptsize $rev$}} & \begin{array}{c} \quad \quad \diagup B \\ \bullet \quad \quad \diagdown A \end{array}
 \end{array} \quad (6.5)$$

$$\begin{array}{ccc}
 \begin{array}{c} \xrightarrow{X} \quad \xrightarrow{X} \\ \bullet \quad \bullet \end{array} & \xrightarrow{\text{\scriptsize $du_X \ du_X$}} & \begin{array}{c} \xrightarrow{X^\sim} \quad \xrightarrow{X^\sim} \\ \bullet \quad \bullet \end{array} \\
 \downarrow g' & & \downarrow g'' \\
 \begin{array}{c} \bullet \quad \bullet \end{array} & \xrightarrow{\text{\scriptsize $rev$}} & \begin{array}{c} \bullet \quad \bullet \end{array}
 \end{array} \quad (6.6)$$

(6.7)

Finally, the morphism

$$rev : a \bullet \rightarrow b \longrightarrow a \bullet \leftarrow b$$

*Notation* . Both morphisms  $rev_a$  and  $du_a$  will be sometimes denoted  $Du_a$ .

Denote  $\mathcal{N}' \subset \mathcal{N}$  the category of nets  $\mathcal{N}_0$  extended by morphisms  $ins_a, del_a$  of insertion and deletion of a 2-vertex with relations (5.1), (5.4), (5.5). There is a desorienting functor  $deso : \mathcal{ON} \rightarrow \mathcal{N}'$  which sends  $rev_a, du_a : \Gamma \rightarrow \Gamma'$  to  $id_{deso(\Gamma)}$ .


**Theorem 6.1.** *The functor  $\text{deso} : \mathcal{ON} \rightarrow \mathcal{N}'$  is an equivalence of categories.*

**6.2. Oriented nets with fusing, twists, braiding and switches.** Now we add to  $\mathcal{ON}$  new morphisms. The obtained category  $ON$  will be made equivalent to  $N$ .

**Definition 6.2.** Let  $\mathcal{ON}$  be a symmetric monoidal category with left cancellations which objects are oriented nets and morphisms are generated over  $\mathcal{ON}$  by the following isomorphisms:

$ins_a : \Gamma \rightarrow \Gamma'$ ,  $del_a : \Gamma' \rightarrow \Gamma$  where  $\Gamma'$  differs from  $\Gamma$  by a 1-vertex  $a$  and an incident edge;


$fus_a : \Gamma \rightarrow \Gamma'$ , where a neighbourhood of  $a$  looks as




(only part of  $\Gamma, \Gamma'$  is drawn here, the complements of it in  $\Gamma$  and  $\Gamma'$  are the same). It is important that cyclic ordering for all 3-vertices is coherent here with the orientation of the plane of drawing;

$Tw_a : \Gamma \rightarrow \Gamma$  for an edge  $a$  of  $\Gamma$ ;

$_a Br_{BC} : \Gamma \rightarrow \Gamma'$  for a 3-vertex  $a \in \Gamma$  and two incident edges  $B, C$  with sources in  $a$ . The net  $\Gamma'$  is obtained from  $\Gamma$  as in unoriented case. Locally the morphism looks as



$S_a : \Gamma \rightarrow \Gamma$  for a loop-edge  $a \in \Gamma$  looking as



They are subject to the following relations. For any glueing  $g : \Gamma_1 \rightarrow \Gamma_2$  whenever upper row in diagrams (5.1), (4.1), (4.5)–(4.7) is defined, the whole diagram is defined and commute. The following diagrams commute

$$\begin{array}{ccc}
 \text{---} \bullet \text{---} & \xrightarrow{ins} & \begin{array}{c} \bullet \\ | \\ \text{---} \bullet \text{---} \end{array} \\
 \parallel & & \downarrow rev \\
 \text{---} \bullet \text{---} & \xrightarrow{ins} & \begin{array}{c} \bullet \\ | \\ \text{---} \bullet \text{---} \end{array}
 \end{array} \quad (6.8)$$

$$\begin{array}{ccccccc}
 \begin{array}{c} a \quad b \\ \text{---} \bullet \text{---} \end{array} & \xrightarrow{Ins} & \begin{array}{c} \bullet \\ | \\ a \quad b \\ \text{---} \bullet \text{---} \end{array} & \xrightarrow{Du_b} & \begin{array}{c} \bullet \\ | \\ a \quad b \\ \text{---} \bullet \text{---} \end{array} & \xrightarrow{Du_a} & \begin{array}{c} \bullet \\ | \\ a \quad b \\ \text{---} \bullet \text{---} \end{array} \\
 \downarrow Del & & & & & & \downarrow Del \\
 \text{---} & \xrightarrow{d \text{ or } rev} & \text{---} & \xrightarrow{Ins} & \text{---} & & \begin{array}{c} a \quad b \\ \text{---} \bullet \text{---} \end{array}
 \end{array} \quad (6.9)$$

$$\begin{array}{ccc}
\begin{array}{c} \xrightarrow{a} \bullet \end{array} & \xrightarrow{ins_b} & \begin{array}{c} \xrightarrow{a} \bullet \xrightarrow{b} \bullet \end{array} \\
ins_c \downarrow & & \downarrow ins_c \\
\begin{array}{c} \xrightarrow{a} \bullet \xrightarrow{c} \bullet \end{array} & \xrightarrow{ins_b} & \begin{array}{c} \xrightarrow{a} \bullet \begin{array}{l} \nearrow c \\ \searrow b \end{array} \end{array}
\end{array} \quad (6.10)$$

$$\begin{array}{ccc}
\begin{array}{c} \xrightarrow{a} \bullet \end{array} & \xrightarrow{ins_b} & \begin{array}{c} \xrightarrow{b} \bullet \xrightarrow{a} \bullet \end{array} \\
ins_c \downarrow & \nearrow \phi & \\
\begin{array}{c} \xrightarrow{a} \bullet \xrightarrow{c} \bullet \end{array} & & 
\end{array} \quad (6.11)$$

(where  $\phi$  is the isomorphism of oriented nets)

$$Tw_a = 1 : \begin{array}{c} \xrightarrow{a} \bullet \end{array} \longrightarrow \begin{array}{c} \bullet \xrightarrow{a} \end{array}. \quad (6.12)$$

and similar diagrams with opposite orientation of edges also commute.

The following diagrams commute

$$\begin{array}{ccccc}
\begin{array}{c} \nearrow \bullet \searrow \end{array} & \xrightarrow{ins} & \begin{array}{c} \nearrow \bullet \xrightarrow{\quad} \bullet \searrow \end{array} & \xrightarrow{ins} & \begin{array}{c} \nearrow \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \searrow \end{array} \\
ins \searrow & & \begin{array}{c} \nearrow \bullet \searrow \end{array} & & \begin{array}{c} \nearrow \bullet \xrightarrow{\quad} \bullet \searrow \end{array} \\
& & \downarrow fus & & \\
& & \begin{array}{c} \nearrow \bullet \xrightarrow{\quad} \bullet \searrow \end{array} & \xrightarrow{ins} & \begin{array}{c} \nearrow \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \searrow \end{array}
\end{array} \quad (6.13)$$

$$\begin{array}{ccccc}
\begin{array}{c} \searrow \bullet \nearrow \end{array} & \xrightarrow{ins} & \begin{array}{c} \searrow \bullet \xrightarrow{\quad} \bullet \nearrow \end{array} & \xrightarrow{ins} & \begin{array}{c} \searrow \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \nearrow \end{array} \\
ins \searrow & & \begin{array}{c} \searrow \bullet \nearrow \end{array} & & \begin{array}{c} \searrow \bullet \xrightarrow{\quad} \bullet \nearrow \end{array} \\
& & \downarrow fus & & \\
& & \begin{array}{c} \searrow \bullet \xrightarrow{\quad} \bullet \nearrow \end{array} & \xrightarrow{ins} & \begin{array}{c} \searrow \bullet \xrightarrow{\quad} \bullet \xrightarrow{\quad} \bullet \nearrow \end{array}
\end{array} \quad (6.14)$$

$$\begin{array}{ccccc}
 \text{crossing} & \xrightarrow{\text{ins}} & \text{horizontal line with dot} & \xrightarrow{\text{ins}} & \text{horizontal line with two dots} \\
 & \searrow \text{ins} & & & \downarrow \text{fus} \\
 & & \text{vertical line with dot} & \xrightarrow{\text{ins}} & \text{vertical line with two dots}
 \end{array} \quad (6.15)$$

$$\begin{array}{ccccccc}
 \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad C \end{array} & \xrightarrow{d_C} & \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad C \end{array} & \xrightarrow{rev} & \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad C \end{array} & \xrightarrow{d_A} & \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad C \end{array} \\
 \downarrow \text{Fus} & & & & & & \downarrow \text{Fus} \\
 \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad C \end{array} & \xrightarrow{d_C} & \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad C \end{array} & \xrightarrow{rev} & \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad C \end{array} & \xrightarrow{d_A} & \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad C \end{array}
 \end{array} \quad (6.16)$$

$$\begin{array}{ccccccc}
 \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad C \end{array} & \xrightarrow{d_A} & \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad C \end{array} & \xrightarrow{rev} & \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad C \end{array} & \xrightarrow{d_D} & \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad C \end{array} \\
 \downarrow \text{Fus} & & & & & & \downarrow \text{Fus} \\
 \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad C \end{array} & \xrightarrow{d_A} & \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad C \end{array} & \xrightarrow{rev} & \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad C \end{array} & \xrightarrow{d_D} & \begin{array}{c} A \quad D \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad C \end{array}
 \end{array} \quad (6.17)$$

$$Tw_A = Tw_B : \begin{array}{c} A \quad B \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \end{array} \longrightarrow \begin{array}{c} A \quad B \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \end{array}, \quad (6.18)$$

$$Tw_a Du_a = Du_a Tw_a \quad (6.20)$$

$$\begin{array}{ccc}
 \begin{array}{c} C \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ A \quad B \end{array} & \xrightarrow{Br_{AB}^{\pm 1}} & \begin{array}{c} C \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad A \end{array} \\
 \downarrow \text{del}_A & & \downarrow \text{del}_A \\
 \begin{array}{c} C \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ A \quad B \end{array} & & \begin{array}{c} C \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ B \quad A \end{array}
 \end{array} \quad (6.21)$$




FIGURE 5. The oriented pentagon equation

$$Br_{BC}^2 = Tw_A Tw_B^{-1} Tw_C^{-1} : \begin{array}{c} A \downarrow \\ \swarrow \quad \searrow \\ B \quad C \end{array} \longrightarrow \begin{array}{c} A \downarrow \\ \swarrow \quad \searrow \\ B \quad C \end{array} \quad (6.22)$$

$$\begin{array}{ccccccc}
 \begin{array}{c} C \downarrow \\ \swarrow \quad \searrow \\ A \quad B \end{array} & \xrightarrow{Br_{AB}} & \begin{array}{c} C \downarrow \\ \swarrow \quad \searrow \\ B \quad A \end{array} & \xrightarrow{du_B} & \begin{array}{c} C \downarrow \\ \swarrow \quad \searrow \\ B^\sim \quad A \end{array} & \xrightarrow{du_C} & \begin{array}{c} C^\sim \downarrow \\ \swarrow \quad \searrow \\ B^\sim \quad A \end{array} \\
 \uparrow Tw_A & & & & & & \downarrow Br_{A,C^\sim} \\
 \begin{array}{c} C \downarrow \\ \swarrow \quad \searrow \\ A \quad B \end{array} & \xleftarrow{rot} & \begin{array}{c} A \downarrow \\ \swarrow \quad \searrow \\ B \quad C \end{array} & \xleftarrow{du_B} & \begin{array}{c} A \downarrow \\ \swarrow \quad \searrow \\ B^\sim \quad C \end{array} & \xleftarrow{du_C} & \begin{array}{c} A \downarrow \\ \swarrow \quad \searrow \\ B^\sim \quad C^\sim \end{array}
 \end{array} \quad (6.23)$$

$$\begin{array}{ccccccc}
\begin{array}{c} C \\ \downarrow \\ A \quad B \end{array} & \xrightarrow{Br_{AB}} & \begin{array}{c} C \\ \downarrow \\ B \quad A \end{array} & \xrightarrow{du_A} & \begin{array}{c} C \\ \downarrow \\ B \quad A^\sim \end{array} & \xrightarrow{du_C} & \begin{array}{c} C^\sim \\ \downarrow \\ B \quad A^\sim \end{array} \\
\uparrow Tw_B & & & & & & \downarrow Br_{C^\sim, B} \\
\begin{array}{c} C \\ \downarrow \\ A \quad B \end{array} & \xleftarrow{rot^{-1}} & \begin{array}{c} B \\ \downarrow \\ C \quad A \end{array} & \xleftarrow{du_A} & \begin{array}{c} B \\ \downarrow \\ C \quad A^\sim \end{array} & \xleftarrow{du_C} & \begin{array}{c} B \\ \downarrow \\ C^\sim \quad A^\sim \end{array}
\end{array} \quad (6.24)$$

$$\begin{array}{ccccc}
\begin{array}{c} X \quad C \\ \diagdown \quad \diagup \\ A \quad B \end{array} & \xrightarrow{Br_{BC}^{\pm 1}} & \begin{array}{c} X \quad B \\ \diagdown \quad \diagup \\ A \quad C \end{array} & \xrightarrow{fus} & \begin{array}{c} X \quad B \\ \diagdown \quad \diagup \\ A \quad C \end{array} \\
\downarrow fus & & & & \downarrow Br_{AC}^{\pm 1} \\
\begin{array}{c} X \quad C \\ \diagdown \quad \diagup \\ A \quad Y \end{array} & \xrightarrow{Br_{YC}^{\pm 1}} & \begin{array}{c} X \quad B \\ \diagdown \quad \diagup \\ C \quad A \end{array} & \xrightarrow{fus} & \begin{array}{c} X \quad B \\ \diagdown \quad \diagup \\ C \quad A \end{array}
\end{array} \quad (6.25)$$

$$(S_M Tw_M)^3 = S_M^2 : \begin{array}{c} \bullet \\ \circlearrowleft \\ \bullet \end{array} \xrightarrow{M} \begin{array}{c} \bullet \\ \circlearrowright \\ \bullet \end{array}, \quad (6.26)$$

$$\begin{array}{ccc}
\begin{array}{c} X \quad M \\ \diagdown \quad \diagup \\ \bullet \end{array} & \xrightarrow{S_M^2} & \begin{array}{c} X \quad M \\ \diagdown \quad \diagup \\ \bullet \end{array} \\
\downarrow {}^X Br_{MM}^{-1} & & \uparrow rev \\
\begin{array}{c} X \quad M \\ \diagdown \quad \diagup \\ \bullet \end{array} & \xrightarrow{Tw_M^{-1}} & \begin{array}{c} X \quad M \\ \diagdown \quad \diagup \\ \bullet \end{array}
\end{array} \quad (6.27)$$

and (6.28) (see Figure 6).


**Theorem 6.2.** *The functor  $deso : ON \rightarrow N$  is an equivalence.*

## 7. Extensions of the category of nets

Let us look for central extensions of the category  $RG$ , or equivalently  $N$ , or  $ON$ . Introduce a new type of morphisms—central charge.

**Definition 7.1.** Let  $C_\Gamma : \Gamma \rightarrow \Gamma$  for each connected oriented net  $\Gamma$  be a morphism commuting with other generators  $du$ ,  $rev$ ,  $ins$ ,  $del$ ,  $fus$ ,  $Tw$ ,  $Br$ ,  $S$ . We assume that

$$C_\emptyset = \text{id}_\emptyset : \emptyset \rightarrow \emptyset. \quad (7.1)$$

FIGURE 6. A relation for  $A_2$  oriented net

If  $\Gamma_1, \Gamma_2, \Gamma$  are connected and  $g : \Gamma_1 \sqcup \Gamma_2 \rightarrow \Gamma$  is a glueing we assume a commutative diagram

$$\begin{array}{ccc}
 \Gamma_1 \sqcup \Gamma_2 & \xrightarrow{C_{\Gamma_1}^k \sqcup C_{\Gamma_2}^l} & \Gamma_1 \sqcup \Gamma_2 \\
 g \downarrow & & \downarrow g \\
 \Gamma & \xrightarrow{C_{\Gamma}^{k+l}} & \Gamma
 \end{array} \tag{7.2}$$

The diagram (7.2) together with (7.1) implies a similar diagram with  $\Gamma_1 \sqcup \dots \sqcup \Gamma_n$ .



We look for extensions of  $ON$  by the generators  $C_\Gamma$  with relations between  $du, rev, fus, Tw, Br, S$  changed by insertion of some powers of  $C_\Gamma$ . We assume that the relations where  $ins, del$  enter are not changed. Then the relations involving  $du, rev, fus, Tw, Br$  also do not change. Indeed, assume that  $C^k$  is inserted into diagrams (6.19)–(6.25) and such, and glue these diagrams with as many  $\bullet \text{---}$  as there are external legs. The generators  $fus, Tw, Br$  will turn into compositions of  $ins$  and  $del$ , and commutativity of such diagram would imply  $k = 0$ . Also the diagram (6.28) after such operation become  $S = SC^k$ , hence, it also does not change.

Therefore, only relations (6.26) and (6.27) could change to

$$(S_M Tw_m)^3 = S_M^2 C^k, \quad (7.3)$$

$$S_M^2 = Br_{MM}^{-1} Tw_M^{-1} rev C^m. \quad (7.4)$$

Renormalizing  $S$  to  $SC^{-k}$  we can restore (7.3). For the sake of symmetry we shall restore (7.4) changing (7.3) though it might require half-integer powers of  $C$ .

**Definition 7.2.** Let  $EN$  be a symmetric monoidal category with left cancellations which objects are oriented nets and morphisms are generated over  $\mathcal{ON}$  by the isomorphisms  $ins, del, fus, Tw, Br, S, C$  with all relations of the category  $ON$  except (6.26) which is substituted by

$$(S_M Tw_M)^3 = C S_M^2 : \rightarrow \bullet \bigcirc \xrightarrow{M} \bullet \bigcirc \xrightarrow{M} \quad (7.5)$$

and relations (7.1), (7.2).

In topology to such extension corresponds the category of framed [24] surfaces. Here we define a central extension  $ESurf \rightarrow Surf$  as a pull-back of the central extension  $EN \rightarrow ON$  along the functor  $Surf \rightarrow RG \rightarrow N \rightarrow ON$  obtained in Theorems 3.1, 5.3, 6.2. Thus the latter extends to a functor  $ESurf \rightarrow EN$ .

## 8. Ribbon categories

*Ribbon* (also *tortile* [20]) category is the following thing: a braided monoidal category  $\mathcal{C}$  [4] with the tensor product  $\otimes$ , the associativity  $a : X \otimes (Y \otimes Z) \rightarrow (X \otimes Y) \otimes Z$ , the braiding (commutativity)  $c : X \otimes Y \rightarrow Y \otimes X$  and a unity object  $I$ , such that  $\mathcal{C}$  is rigid (for any object  $X \in \mathcal{C}$  there are dual objects  $\smile X$  and  $X \smile$  with evaluations  $ev : \smile X \otimes X \rightarrow I$ ,  $ev : X \otimes X \smile \rightarrow I$  and coevaluations  $coev : I \rightarrow X \otimes \smile X$ ,

$\text{coev} : I \rightarrow X^\sim \otimes X$ ) and possess a ribbon twist  $\nu$ . A ribbon twist [4, 17, 20]  $\nu = \nu_X : X \rightarrow X$  is a self-adjoint ( $\nu_{X^\sim} = \nu_X^t$ ) functorial automorphism such that  $c^2 = \nu_X^{-1} \otimes \nu_Y^{-1} \circ \nu_{X \otimes Y}$ . In a ribbon category there are functorial isomorphisms [9]

$$u_1^2 = \begin{array}{c} X \\ \diagdown \\ \text{---} \circ \text{---} \\ \diagup \\ X^{\sim\sim} \end{array}, \quad u_{-1}^2 = \begin{array}{c} X \\ \diagdown \\ \text{---} \circ \text{---} \\ \diagup \\ X^{\sim\sim} \end{array}, \quad u_1^{-2} = \begin{array}{c} X \\ \diagup \\ \text{---} \circ \text{---} \\ \diagdown \\ {}^{\sim\sim}X \end{array}, \quad u_{-1}^{-2} = \begin{array}{c} X \\ \diagup \\ \text{---} \circ \text{---} \\ \diagdown \\ {}^{\sim\sim}X \end{array}$$

$$u_0^2 = u_1^2 \circ \nu^{-1} = u_{-1}^2 \circ \nu : X \rightarrow X^{\sim\sim}, \quad u_0^{-2} = u_1^{-2} \circ \nu^{-1} = u_{-1}^{-2} \circ \nu : X \rightarrow {}^{\sim\sim}X.$$

Changing the category  $\mathcal{C}$  by an equivalent one, we can (and we will) assume that  ${}^{\sim}X = X^\sim$ ,  $X^{\sim\sim} = {}^{\sim\sim}X = X$  and  $u_0^2 = u_0^{-2} = \text{id}_X$  (see [9]).

If in addition  $\mathcal{C}$  is additive, it is  $k$ -linear with  $k = \text{End } I$ . We assume in the following that  $k$  is a field, in which each element has a square root. Often  $\mathcal{C}$  will be noetherian abelian category with finite dimensional  $k$ -vector spaces  $\text{Hom}_{\mathcal{C}}(A, B)$ . In such case there exists a coend  $F = \int X \otimes X^\sim$  as an object of a cocompletion  $\hat{\mathcal{C}}$  [9] of  $\mathcal{C}$ . It is a Hopf algebra (see [10, 12, 15]). There is a Hopf pairing  $\omega : F \otimes F \rightarrow I$  [10],

$$\omega = \begin{array}{c} F \quad \quad F \\ \diagdown \quad \diagup \\ \text{---} \quad \text{---} \\ \diagup \quad \diagdown \\ \text{---} \quad \text{---} \end{array}$$

such that

$$\text{Ker } \omega = \text{Ker}^{\text{left}} \omega = \text{Ker}^{\text{right}} \omega \in \hat{\mathcal{C}}.$$

The quotient  $\mathbf{f} = F / \text{Ker } \omega \in \hat{\mathcal{C}}$  is also a Hopf algebra and the first modular axiom is [10]

(M1)  $\mathbf{f}$  is an object of  $\mathcal{C}$  (and not only of a cocompletion  $\hat{\mathcal{C}}$ ) (more scrupulously, it means that there exists an exact sequence  $0 \rightarrow \text{Ker } \omega \rightarrow F \rightarrow \mathbf{f} \rightarrow 0$  in  $\hat{\mathcal{C}}$ , where  $\mathbf{f}$  is an object from  $\mathcal{C} \subset \hat{\mathcal{C}}$ ).

Being the coend  $\int X \otimes X^\sim$ , the object  $F \in \hat{\mathcal{C}}$  has an automorphism  $\underline{\nu \otimes 1} \stackrel{\text{def}}{=} \int \nu \otimes 1$  (notations are from [10]). The second modular axiom says [10]

$$(M2) \quad \underline{\nu \otimes 1}(\text{Ker } \omega) \subset \text{Ker } \omega$$

(more scrupulously, there exist morphisms  $T' : \text{Ker } \omega \rightarrow \text{Ker } \omega \in \hat{\mathcal{C}}$ ,  $T : \mathbf{f} \rightarrow \mathbf{f} \in \mathcal{C}$  such that the diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & \text{Ker } \omega & \rightarrow & F & \rightarrow & \mathbf{f} \rightarrow 0 \\ & & T' \downarrow & & \underline{\nu \otimes 1} \downarrow & & T \downarrow \\ 0 & \rightarrow & \text{Ker } \omega & \rightarrow & \bar{F} & \rightarrow & \mathbf{f} \rightarrow 0 \end{array}$$

commutes).

**Definition 8.1.** A noetherian abelian ribbon category  $\mathcal{C}$  with finite dimensional  $k$ -vector spaces of morphisms  $\text{Hom}_{\mathcal{C}}(A, B)$  is called *modular*, if axioms (M1), (M2) are satisfied.

It was shown in [10] that in the case of a modular category there exists a morphism  $\mu : I \rightarrow \mathbf{f}$ , which is the integral of a dual Hopf algebra  $\check{\mathbf{f}} \simeq \mathbf{f}$ , and

$$\nu^{-1} \left[ \text{cup with } \mu \text{ on top} \right] = \lambda^{-1} \left[ \text{vertical line with } \nu \text{ on right} \right], \quad \nu \left[ \text{cup with } \mu \text{ on top} \right] = \lambda \left[ \text{vertical line with } \nu^{-1} \text{ on right} \right]$$

for some invertible constant  $\lambda \in k^\times$ . The pair  $(\mu, \lambda)$  is unique up to a sign. Morphisms  $S, S^{-1} : \mathbf{f} \rightarrow \mathbf{f}$

$$S = \left[ \text{cup with } \mu \text{ on top} \right], \quad S^{-1} = \left[ \text{cap with } \mu \text{ on top} \right]$$

are inverse to each other. Morphisms  $S$  and  $T$  (defined via (M2)) yield a projective representation of a mapping class group of a torus with one hole:

$$(ST)^3 = \lambda S^2, \quad S^2 = \gamma^{-1},$$

$$T\gamma = \gamma T, \quad \gamma^2 = \nu.$$

Here  $\gamma : \mathbf{f} \rightarrow \mathbf{f}$  is the antipode of the Hopf algebra  $\mathbf{f}$ ,

$$\gamma = \begin{array}{c} \mathbf{f} \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ \mathbf{f} \end{array} \bigg/ \bigcirc$$

## 9. From a ribbon category to a modular functor

Assume that we are given a small ribbon category  $(\mathcal{C}, \otimes, a, c, \nu, I)$  for which the duality map  $\cdot^\sim : \text{Ob } \mathcal{C} \rightarrow \text{Ob } \mathcal{C}$  is involutive and  $u_0^2 : A \rightarrow A^\sim = A$  equals  $1_A$  for all objects  $A \in \mathcal{C}$ . Further, we assume that  $\mathcal{C}$  is a noetherian abelian category and  $\text{End } I = k$  is a field, thus,  $\mathcal{C}$  is  $k$ -linear. We assume also that  $k$ -vector spaces  $\text{Hom}(A, B)$  are finite dimensional for any  $A, B \in \mathcal{C}$ . Finally, we assume that algebra  $\mathbf{f}$ , quotient of  $F$ , belongs to  $\mathcal{C}$  and  $\text{Ker}(F \rightarrow \mathbf{f}) = \text{Ker } \omega$  is  $\underline{\nu \otimes 1}$ -invariant.

### 9.1. Preliminaries about exact functors.

**Lemma 9.1.** *Let  $F : \mathcal{C} \rightarrow k\text{-Vect}$ ,  $G : \mathcal{C}^{op} \rightarrow k\text{-Vect}$  be functors. Then*

$$\begin{aligned} \int^X F(X) \otimes \text{Hom}(X, B) &\rightarrow F(B), & v \otimes f &\mapsto F(f).v \\ \int^X \text{Hom}(B, X) \otimes G(X) &\rightarrow G(B), & f \otimes v &\mapsto G(f).v \end{aligned}$$

*are isomorphisms of vector spaces.*

Let  $F : (\mathcal{C}^{k+n})^{op} \times \mathcal{C}^{n+l} \rightarrow k\text{-vect}$  be a  $k$ -linear left exact (preserving kernels) functor. By the “parameter theorem for coends” of Mac Lane [13] a coend of the bifunctor

$$F(A, -; -; B) : (\mathcal{C}^n)^{op} \times \mathcal{C}^n \rightarrow k\text{-vect}$$

with fixed  $A \in \mathcal{C}^k$ ,  $B \in \mathcal{C}^l$  is identified as a functor in  $A, B$  with a coend of the bifunctor

$$F' : (\mathcal{C}^n)^{op} \times \mathcal{C}^n \rightarrow \text{Functors}((\mathcal{C}^{op})^k \times \mathcal{C}^l \rightarrow k\text{-vect}).$$

Here functors are  $k$ -linear functors. But, generally, the coend of the bifunctor

$$F'' : (\mathcal{C}^n)^{op} \times \mathcal{C}^n \rightarrow \text{Left Exact Functors}((\mathcal{C}^{op})^k \times \mathcal{C}^l \rightarrow k\text{-vect}).$$

is a different thing. In any case we have a morphism

$$\int^X F'(X, X) \rightarrow \int^X F''(X, X).$$

To stress the difference, we introduce different notations for the usual coend

$$\int^X F(A, X; X, B) \equiv \left( \int^X F'(X, X) \right) (A, B)$$

and for the second type of coend, the left exact functor

$$\oint^X F(A, X; X, B) \stackrel{\text{def}}{=} \left( \int^X F''(X, X) \right) (A, B).$$

The Fubini theorem for left exact functors (repeated coend theorem) is proved exactly in the same way as for arbitrary functors (see Mac Lane's book [13]). Nevertheless, we present a proof here for the sake of completeness.

**Theorem 9.2.** *Let  $\mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_3, \mathcal{A}$  be abelian  $k$ -linear categories and let  $F : \mathcal{C}_2^{op} \times \mathcal{C}_1^{op} \times \mathcal{C}_1 \times \mathcal{C}_2 \times \mathcal{C}_3 \rightarrow \mathcal{A}$  be left exact  $k$ -linear functor. Assume that there exists coend with parameters*

$$j_X(U; V, A) : F(U, X; X, V, A) \rightarrow G(U; V, A) = \int^{X \in \mathcal{C}_1} F(U, X; X, V, A).$$

*Assume that  $G$ , viewed as a functor  $G : \mathcal{C}_2^{op} \times \mathcal{C}_2 \rightarrow \text{Functors}(\mathcal{C}_3 \rightarrow \mathcal{A})$ , has a left exact coend  $H \in \text{LEF}(\mathcal{C}_3 \rightarrow \mathcal{A})$  (set of Left Exact Functors)*

$$k_U(A) : G(U; U, A) \rightarrow H(A) = \int^U G(U; U, A)$$

*(that means that  $k$  is dinatural,  $H$  is left exact and the pair  $(k, H)$  is universal between all dinatural transformations of  $G$  into left exact functors). Then*

$$i_{X,U}(A) : F(U, X; X, U, A) \xrightarrow{j_X(U;U,A)} G(U; U, A) \xrightarrow{k_U(A)} H(A)$$

*is a coend  $\oint^{X,U} F(U, X; X, U, A)$  of  $F$ , viewed as a bifunctor*

$$(\mathcal{C}_1 \times \mathcal{C}_2)^{op} \times (\mathcal{C}_1 \times \mathcal{C}_2) \rightarrow \text{LEF}(\mathcal{C}_3 \rightarrow \mathcal{A}).$$

We write the statement of the theorem as an isomorphism

$$\oint^{X,U} F(U, X; X, U, A) \simeq \oint^U \int^X F(U, X; X, U, A).$$

In particular case  $\mathcal{C}_2 = 0$  we have

**Corollary 9.3.** *Let  $G(A) = \int^{X \in \mathcal{C}_1} F(X; X, A)$  for left exact  $k$ -linear functor  $F : \mathcal{C}_1^{op} \times \mathcal{C}_1 \times \mathcal{C}_3 \rightarrow \mathcal{A}$ . Then*

$$\oint^X F(X; X, A) = \oint^0 G(A)$$

if there exists a left exact functor  $\oint^0 G : \mathcal{C}_3 \rightarrow \mathcal{A}$  with isomorphisms  $\text{Nat}(G, E) \cong \text{Nat}(\oint^0 G, E)$  for any left exact  $E : \mathcal{C}_3 \rightarrow \mathcal{A}$ .

**9.2. Modular functor.** Our goal is to construct a functor  $Z : EN \rightarrow k\text{-vect}$ , which will satisfy to the following conditions:

- (i)  $Z$  is a symmetric monoidal functor.
- (ii) To  $D_2$  corresponds

$$Z(D_2; B; C) = \text{Hom}(B, C),$$

to glueing of such corresponds the composition

$$\text{Hom}(A, B) \otimes \text{Hom}(B, C) \rightarrow \text{Hom}(A, C).$$

- (iii) To  $D_3$  correspond

$$Z \left( \begin{array}{c} A \quad B \\ \swarrow \quad \searrow \\ \bullet \\ \downarrow \\ C \end{array} \right) = \text{Hom}(A \otimes B, C),$$

$$Z \left( \begin{array}{c} A \\ \downarrow \\ \bullet \\ \swarrow \quad \searrow \\ B \quad C \end{array} \right) = \text{Hom}(A, B \otimes C).$$

To glueing of  $D_3$  with  $D_2$  corresponds an action of  $\text{Hom}(-, -)$  on  $\text{Hom}(- \otimes -, -)$  or  $\text{Hom}(-, - \otimes -)$  via composition.

- (iv) To  $D_1$  correspond

$$Z(D_1; X; ) = \text{Hom}(X, I),$$

$$Z(D_1; ; X) = \text{Hom}(I, X),$$

where  $I$  is unity object of  $\mathcal{C}$ . To glueing of  $D_1$  with  $D_2$  correspond compositions

$$\text{Hom}(X, Y) \otimes \text{Hom}(Y, I) \rightarrow \text{Hom}(X, I),$$

$$\text{Hom}(I, X) \otimes \text{Hom}(X, Y) \rightarrow \text{Hom}(I, Y).$$

- (v) To isomorphisms

$$du_B : \begin{array}{c} A \quad B \\ \swarrow \quad \searrow \\ \bullet \\ \downarrow \\ C \end{array} \longrightarrow \begin{array}{c} A \quad B^\sim \\ \swarrow \quad \searrow \\ \bullet \\ \downarrow \\ C \end{array},$$

$$du_A : \begin{array}{c} A \quad B \\ \swarrow \quad \searrow \\ \bullet \\ \downarrow \\ C \end{array} \longrightarrow \begin{array}{c} A^\sim \quad B \\ \swarrow \quad \searrow \\ \bullet \\ \downarrow \\ C \end{array}$$

correspond adjunctions

$$d_l : \text{Hom}(A \otimes B, C) \rightarrow \text{Hom}(A, C \otimes B^\sim),$$

$$d_r : \text{Hom}(A \otimes B, C) \rightarrow \text{Hom}(B, A^\sim \otimes C).$$

(vi) For any oriented net  $\Gamma$  glueings with  $D_2$  makes  $Z(\Gamma)$  into a functor

$$Z(\Gamma) : (\mathcal{C}^{op})^k \times \mathcal{C}^l \rightarrow k\text{-vect}$$

We assume that  $Z(\Gamma)$  is left exact (additive and preserving kernels).

(vii) If  $f : \Gamma \rightarrow \tilde{\Gamma}$  is a glueing and  $g(\Gamma) = g(\tilde{\Gamma})$ , then the morphism of functors

$$\int^{X \in \mathcal{C}^k} Z(\Gamma, \dots, X; X, \dots) \rightarrow Z(\tilde{\Gamma}; \dots; \dots)$$

is an isomorphism.

(viii) To the morphism

The diagram shows a morphism labeled  $fus$ . On the left, there are two separate objects: the top one has inputs  $X$  and  $C$  and output  $A$ ; the bottom one has inputs  $A$  and  $B$  and output  $C$ . An arrow points from the output of the top object to the input of the bottom object. On the right, the result is a single object with inputs  $X$  and  $C$  and output  $B$ . A vertical line labeled  $V$  connects the two input nodes.

corresponds

$$\begin{aligned} \text{Hom}(X, A \otimes (B \otimes C)) &\simeq \int^{U \in \mathcal{C}} \text{Hom}(X, A \otimes U) \otimes \text{Hom}(U, B \otimes C) \simeq \\ &\simeq Z \left( \begin{array}{c} X \quad C \\ \swarrow \quad \searrow \\ \bullet \quad \bullet \\ \swarrow \quad \searrow \\ A \quad B \end{array} \right) \xrightarrow{Z(fus)} Z \left( \begin{array}{c} X \quad C \\ \swarrow \quad \searrow \\ \bullet \quad \bullet \\ \swarrow \quad \searrow \\ A \quad B \end{array} \right) \simeq \\ &\simeq \int^{V \in \mathcal{C}} \text{Hom}(X, V \otimes C) \otimes \text{Hom}(V, A \otimes B) \simeq \text{Hom}(X, (A \otimes B) \otimes C), \end{aligned}$$

which coincides with  $\text{Hom}(X, a_{A,B,C})$ .

(ix) To the morphism

The diagram shows a morphism labeled  $Br_{AB}$ . On the left, there is an object with inputs  $A$  and  $B$  and output  $X$ . On the right, there is an object with inputs  $B$  and  $A$  and output  $X$ . An arrow points from the left object to the right object.

corresponds

$$\text{Hom}(X, c_{A,B}) : \text{Hom}(X, A \otimes B) \rightarrow \text{Hom}(X, B \otimes A).$$

(x) To the morphism

The diagram shows a morphism labeled  $Tw_Y$ . On the left, there are two objects: the top one has input  $X$  and output  $Y$ ; the bottom one has input  $Y$  and output  $X$ . An arrow points from the output of the top object to the input of the bottom object. On the right, the result is a single object with input  $X$  and output  $Y$ .

corresponds

$$\text{Hom}(X, \nu_Y) : \text{Hom}(X, Y) \rightarrow \text{Hom}(X, Y).$$

(xi) To a tadpole  $A_1$ , obtained via glueing from  $D_3$ , corresponds

$$Z \left( \begin{array}{c} X \quad \nearrow M \\ \leftarrow \quad \bullet \quad \searrow \\ M \end{array} \right) = \text{Hom}(X \otimes M, M) \simeq \text{Hom}(X, M \otimes M) \rightarrow$$

$$\rightarrow \text{Hom}(X, \mathbf{f}) = Z \left( \begin{array}{c} X \quad \circlearrowright M \end{array} \right).$$

$Z$  can be viewed as a monoidal functor

$$\{\text{extended nets}\} \rightarrow \hat{\mathcal{C}}_{*,*}.$$

Now we shall construct such a functor step by step, proving the following

**Theorem 9.4.** *There exists one and only one up to equivalence functor  $Z : EN \rightarrow k\text{-vect}$ , satisfying assumptions (i)–(xi) above. It has also property:*

(xii) *Let  $f : \Gamma \rightarrow \tilde{\Gamma}$  be a glueing and let the boundary of each connected component of  $\tilde{\Gamma}$  not be empty. The functor*

$$Z(\Gamma; \dots, X; X, \dots) : (\mathcal{C}^{op})^{k+n} \times \mathcal{C}^{n+l} \rightarrow k\text{-vect}$$

*can be represented as a bifunctor*

$$Z'(\Gamma)(X, X) : (\mathcal{C}^n)^{op} \times (\mathcal{C}^n) \rightarrow \hat{\mathcal{C}}_{k,l},$$

*where  $\hat{\mathcal{C}}_{k,l}$  is the category of left exact functors  $(\mathcal{C}^{op})^k \times \mathcal{C}^l \rightarrow k\text{-Vect}$ . The coend of this bifunctor is mapped to  $Z(\tilde{\Gamma}) \in \hat{\mathcal{C}}_{k,l}$*

$$\int^{X \in \mathcal{C}^k} Z'(\Gamma)(X, X) \rightarrow Z(\tilde{\Gamma}).$$

*We claim that this is an epimorphism in  $\hat{\mathcal{C}}_{k,l}$ .*

Proof is the matter of the remaining part of this chapter.

**9.3. A functor on the category of oriented nets.** Let  $\mathcal{ON}_>$  be a symmetric monoidal category of glueings of oriented nets, having at least one end at each connected component. Let us construct a functor  $\mathcal{Z} : \mathcal{ON}_> \rightarrow k\text{-vect}$ . Fix the value of the functor  $\mathcal{Z}$  on elementary objects— $D_1, D_2, D_3$  as in (ii)–(iv). So it is fixed on disjoint unions  $\bigsqcup_i X_i$  of such objects:  $\mathcal{Z}(\bigsqcup_i X_i) = \otimes_i \mathcal{Z}(X_i)$ . (In fact, it is fixed only up to a permutation of tensor multiplicands. To fix it completely, we could define nets with chosen total ordering of the set  $V_3 \sqcup V_2 \sqcup V_1$ , and add to  $\mathcal{ON}$  new morphisms, which change only that ordering. Obtained category is equivalent to the old one. We shall not remind about such tricks in the following.)



The following condition determines the value of  $\mathcal{Z}$  on  $\mathcal{ON}_>$ :

(xiii) Let  $f : \Gamma \rightarrow \tilde{\Gamma}$  be a glueing, and let the boundary of each connected component of  $\tilde{\Gamma}$  be not empty, then there exists a coend  $\oint \mathcal{Z}(\Gamma)$  and the morphism

$$\oint^{X \in \mathcal{C}^k} \mathcal{Z}(\Gamma; \dots, X; X, \dots) \rightarrow \mathcal{Z}(\tilde{\Gamma}; \dots; \dots)$$

is an isomorphism of left exact functors.

Let  $\Gamma \in \mathcal{ON}_>$  be a connected net of genus  $g$  with  $k$  incoming and  $l$  outgoing legs,  $k + l > 0$ . We construct a space  $\mathcal{Z}(\Gamma)$  in the following way. Let  $\Gamma_1$  be the net obtained from  $\Gamma$  by cutting all internal edges. There is a canonical glueing  $f : \Gamma_1 \rightarrow \Gamma$ . We show the existence of the coend

$$\oint^{X \in \mathcal{C}^n} \mathcal{Z}(\Gamma_1; A, X; X, B) \in \hat{\mathcal{C}}_{k,l}$$

and define  $\mathcal{Z}(\Gamma)$  to be that functor.

Cut  $\Gamma$  at such edges  $w_1, \dots, w_g$  that the obtained net  $\Gamma_2$  is a tree. Introduce “internal variables”  $W_1, \dots, W_g \in \mathcal{C}$  corresponding to  $w_i$ ,  $1 \leq i \leq g$ , and let  $Y_1, \dots, Y_{n-g} \in \mathcal{C}$  be other “internal variables” corresponding to other edges. Cutting along them breaks  $\Gamma_2$  into pieces, forming  $\Gamma_1$ . We have the whole collection of variables  $\{X_1, \dots, X_n\} = \{W_1, \dots, W_g, Y_1, \dots, Y_{n-g}\}$ . We apply Theorem 9.2. The coend  $\int^{Y \in \mathcal{C}^{n-g}} \mathcal{Z}(\Gamma_1; A, W, Y; Y, V, B)$  exists and gives the functor  $\mathcal{Z}(\Gamma_2; A, W; V, B)$ . It is isomorphic to a functor

$$\text{Hom}(A_1 \otimes \dots \otimes A_k, B_1 \otimes \dots \otimes B_l \otimes (V_1 \otimes W_1^\vee) \otimes \dots \otimes (V_g \otimes W_g^\vee))$$

with some parenthesis for tensor product. This is proved by induction on the number of vertices of the tree. Inductive step use Lemma 9.1 similarly to calculation

$$\begin{aligned} & \int^X \text{Hom}(A, X \otimes B) \otimes \text{Hom}((C \otimes X) \otimes D), E) \stackrel{d_B}{\simeq} \\ & \simeq \int^X \text{Hom}(A \otimes B^\vee, X) \otimes \text{Hom}((C \otimes X) \otimes D), E) \simeq \text{Hom}((C \otimes (A \otimes B^\vee)) \otimes D, E). \end{aligned}$$

By Theorem 9.2

$$\begin{aligned} & \oint^{X \in \mathcal{C}^n} \mathcal{Z}(\Gamma_1; A, X; X, B) \simeq \oint^{W \in \mathcal{C}^g} \mathcal{Z}(\Gamma_2; A, W; W; B) \simeq \\ & \simeq \oint^{W \in \mathcal{C}^g} \text{Hom}(A_1 \otimes \dots \otimes A_k, B_1 \otimes \dots \otimes B_l \otimes (W_1 \otimes W_1^\vee) \otimes \dots \otimes (W_g \otimes W_g^\vee)) \end{aligned}$$

if the latter exists. We show that, indeed, it exists and equals

$$\mathrm{Hom}(A_1 \otimes \dots \otimes A_k, B_1 \otimes \dots \otimes B_l \otimes (\int^{W_1 \in \mathcal{C}} W_1 \otimes W_1^\sim) \otimes \dots \otimes (\int^{W_g \in \mathcal{C}} W_g \otimes W_g^\sim))$$

for  $k + l > 0$ .

Using duality isomorphisms, we can assume that  $k = 1$ . Assume that we are given functorial morphisms

$$i_{W_1 \dots W_g}(A, B) : \mathrm{Hom}(A, B_1 \otimes \dots \otimes B_l \otimes C \otimes (W_1 \otimes W_1^\sim) \otimes \dots \otimes (W_g \otimes W_g^\sim)) \rightarrow G(A; B_1, \dots, B_l)$$

which define dinatural transformation. Here  $C$  is an object of  $\hat{\mathcal{C}}$  and  $G : \mathcal{C}^{op} \times \mathcal{C}^l \rightarrow k\text{-Vect}$  is a left exact functor. We prove by induction on  $g$  that they all factorize through the unique morphism

$$\mathrm{Hom}(A, B_1 \otimes \dots \otimes B_l \otimes C \otimes (\int^{W_1} W_1 \otimes W_1^\sim) \otimes \dots \otimes (\int^{W_g} W_g \otimes W_g^\sim)) \rightarrow G(A; B).$$

If  $g \geq 1$  denote by  $D$  the product  $B_1 \otimes \dots \otimes B_l \otimes C \otimes (W_1 \otimes W_1^\sim) \otimes \dots \otimes (W_{g-1} \otimes W_{g-1}^\sim)$  and denote by  $G_1(A)$  the left exact functor  $G(A; B_1, \dots, B_l)$  with fixed  $B_1, \dots, B_l$ . There exists tautologically  $T \in \hat{\mathcal{C}}$ , such that  $G_1(A) = \mathrm{Hom}(A, T)$ . Dinaturality implies commutativity of the diagram

$$\begin{array}{ccc} \mathrm{Hom}(A, D \otimes (V_g \otimes W_g^\sim)) & \xrightarrow{\mathrm{Hom}(A, D \otimes f \otimes W_g^\sim)} & \mathrm{Hom}(A, D \otimes (W_g \otimes W_g^\sim)) \\ \mathrm{Hom}(A, D \otimes (V_g \otimes f^t)) \downarrow & & \downarrow i_{\dots, W_g} \\ \mathrm{Hom}(A, D \otimes (V_g \otimes V_g^\sim)) & \xrightarrow{i_{\dots, V_g}} & \mathrm{Hom}(A, T) \end{array}$$

Hence, the following diagram in  $\hat{\mathcal{C}}$  commutes for any  $f : V_g \rightarrow W_g$ :

$$\begin{array}{ccc} D \otimes (V_g \otimes W_g^\sim) & \xrightarrow{D \otimes f \otimes W_g^\sim} & D \otimes (W_g \otimes W_g^\sim) \\ D \otimes (V_g \otimes f^t) \downarrow & & \downarrow \\ D \otimes (V_g \otimes V_g^\sim) & \longrightarrow & T \end{array}$$

Consequently, there is a morphism in  $\hat{\mathcal{C}}$

$$D \otimes \int^W W \otimes W^\sim \cong \int^W D \otimes (W \otimes W^\sim) \rightarrow T.$$

The first isomorphism here follows from the fact that  $\otimes$  preserves colimits in  $\hat{\mathcal{C}}$ . Thus, for any collection of objects  $B_1, \dots, B_l, W_1, \dots, W_{g-1}$

we obtained morphisms

$$j_{B_1, \dots, B_l, W_1, \dots, W_{g-1}}(A) : \text{Hom}(A, B_1 \otimes \dots \otimes B_l \otimes C \otimes (W_1 \otimes W_1^\sim) \otimes \dots \otimes (W_{g-1} \otimes W_{g-1}^\sim) \otimes (\int^{W_g} W_g \otimes W_g^\sim)) \rightarrow G(A; B_1, \dots, B_l)$$

functorial in  $A$ . Morphisms  $i_W(A; B)$  are factorized through  $j_{B,W}(A)$  and the latter are characterized by that property.

One can prove that  $j_W(A; B) \equiv j_{B,W}(A)$  is functorial in  $A$  and  $B_i$  and that  $j_W$  is dinatural in  $W$ .

Constant tensor multiple  $F = \int^{W_p} W_p \otimes W_p^\sim$  can be adjoint to  $C$ . We finish computation of  $\oint \mathcal{Z}(\Gamma_1)$  by induction.

Thus, we defined a functor  $\mathcal{Z} : \mathcal{ON} \rightarrow k\text{-Vect}$ .

$$\mathcal{Z}(\Gamma) \stackrel{\text{def}}{=} \oint \mathcal{Z}(\Gamma_1) \simeq \text{Hom}(A_1 \otimes \dots \otimes A_k, B_1 \otimes \dots \otimes B_l \otimes F \otimes \dots \otimes F).$$

This functor satisfies (i)–(iv), (vi), (vii).

Now we prove the property (xiii) for  $\mathcal{Z}$ . We assume that  $\tilde{\Gamma}$  is connected with non-empty boundary. Any glueing  $f : \Gamma \rightarrow \tilde{\Gamma}$  can be factorized into  $\Gamma \xrightarrow{f_1} \Gamma' \xrightarrow{f_2} \tilde{\Gamma}$ , where genus of  $\Gamma'$  equals genus of  $\Gamma$  and  $\Gamma'$  is connected. Let variables  $X_i$  correspond to circles, glued by  $f_1$ , and let  $Y_j$  correspond to ones, glued by  $f_2$ . Considered morphism  $\mathcal{Z}(f)$  factorizes as

$$\oint \mathcal{Z}(\Gamma) \simeq \oint^Y \int^X \mathcal{Z}(\Gamma) \simeq \oint^Y \mathcal{Z}(\Gamma') \rightarrow \mathcal{Z}(\Gamma)$$

by Theorem 9.2 and (vii). This is an isomorphism, because

$$\begin{aligned} \oint^Y \text{Hom}(A_1 \otimes \dots \otimes A_k, B_1 \otimes \dots \otimes B_l \otimes (Y_1 \otimes Y_1^\sim) \otimes \dots \otimes (Y_p \otimes Y_p^\sim) \otimes F \otimes \dots \otimes F) \rightarrow \\ \rightarrow \text{Hom}(A_1 \otimes \dots \otimes A_k, B_1 \otimes \dots \otimes B_l \otimes F \otimes \dots \otimes F \otimes \dots \otimes F) \end{aligned}$$

is an isomorphism.

**9.4. Relations for insertions and reversals.** We represent morphisms  $X = \text{ins}, \text{del} : \Gamma \rightarrow \Gamma'$  from Section 6 by the following procedure. Let  $\Gamma = \Gamma_1 \cup \sigma$ ,  $\Gamma' = \Gamma_1 \cup \sigma'$ , where  $X = \text{ins}, \text{del} : \sigma \rightarrow \sigma'$  is a standard morphism, and  $X|_{\Gamma_1} = \text{id}$ . Then we define  $\mathcal{Z}(X)$  as

$$\mathcal{Z}(\Gamma) \simeq \oint \mathcal{Z}(\Gamma_1) \otimes \mathcal{Z}(\sigma) \xrightarrow{\oint \text{id} \otimes \mathcal{Z}(X)} \oint \mathcal{Z}(\Gamma_1) \otimes \mathcal{Z}(\sigma') \simeq \mathcal{Z}(\Gamma'),$$

an isomorphism of coends, induced by isomorphism of underlying bifunctors.

Now we construct a functor  $\mathcal{Z}$  on  $\mathcal{ON}_>$  extended by *ins* and *del* satisfying the axioms. To deletion or insertion of a 2-vertex correspond isomorphisms of Lemma 9.1. To deletion or insertion of a 1-vertex correspond isomorphisms which are glueings of identity with

$$\begin{aligned}\mathcal{Z}\left(\begin{array}{c} X \downarrow \\ \hline \rightarrow \bullet \rightarrow Y \end{array} \longrightarrow \begin{array}{c} X \quad Y \\ \hline \rightarrow \bullet \rightarrow \end{array}\right) &= \text{Hom}(X \otimes I, Y) \xrightarrow{\text{Hom}(r_X^{-1}, Y)} \text{Hom}(X, Y) \\ \mathcal{Z}\left(\begin{array}{c} Y \downarrow \\ \hline \rightarrow \bullet \rightarrow X \end{array} \longrightarrow \begin{array}{c} Y \quad X \\ \hline \rightarrow \bullet \rightarrow \end{array}\right) &= \text{Hom}(I \otimes X, Y) \xrightarrow{\text{Hom}(l_X^{-1}, X)} \text{Hom}(X, Y) \\ \mathcal{Z}\left(\begin{array}{c} X \downarrow \\ \hline \rightarrow \bullet \rightarrow Y \end{array} \longrightarrow \begin{array}{c} X \quad Y \\ \hline \rightarrow \bullet \rightarrow \end{array}\right) &= \text{Hom}(X, Y \otimes I) \xrightarrow{\text{Hom}(X, r_Y)} \text{Hom}(X, Y) \\ \mathcal{Z}\left(\begin{array}{c} Y \downarrow \\ \hline \rightarrow \bullet \rightarrow X \end{array} \longrightarrow \begin{array}{c} Y \quad X \\ \hline \rightarrow \bullet \rightarrow \end{array}\right) &= \text{Hom}(X, I \otimes Y) \xrightarrow{\text{Hom}(X, l_Y)} \text{Hom}(X, Y)\end{aligned}$$

The orientation reversing morphism  $du$  is realized on external legs as a glueing of identity with duality adjunctions

$$\begin{aligned}\mathcal{Z}(du_A) : \mathcal{Z}\left(\begin{array}{c} X \downarrow \\ \swarrow \quad \searrow \\ A \quad B \end{array}\right) &= \text{Hom}(X, A \otimes B) \xrightarrow{d_r^{-1}} \text{Hom}(A^\sim \otimes X, B) = \mathcal{Z}\left(\begin{array}{c} X \downarrow \\ \swarrow \quad \searrow \\ A^\sim \quad B \end{array}\right), \\ \mathcal{Z}(du_B) : \mathcal{Z}\left(\begin{array}{c} X \downarrow \\ \swarrow \quad \searrow \\ A \quad B \end{array}\right) &= \text{Hom}(X, A \otimes B) \xrightarrow{d_l^{-1}} \text{Hom}(X \otimes B^\sim, A) = \mathcal{Z}\left(\begin{array}{c} X \downarrow \\ \swarrow \quad \searrow \\ A \quad B^\sim \end{array}\right), \\ \mathcal{Z}(du_X) : \mathcal{Z}\left(\begin{array}{c} X \quad Y \\ \swarrow \quad \searrow \\ A \downarrow \end{array}\right) &= \text{Hom}(X \otimes Y, A) \xrightarrow{d_r} \text{Hom}(Y, X^\sim \otimes A) = \mathcal{Z}\left(\begin{array}{c} X^\sim \quad Y \\ \swarrow \quad \searrow \\ A \downarrow \end{array}\right), \\ \mathcal{Z}(du_Y) : \mathcal{Z}\left(\begin{array}{c} X \quad Y \\ \swarrow \quad \searrow \\ A \downarrow \end{array}\right) &= \text{Hom}(X \otimes Y, A) \xrightarrow{d_l} \text{Hom}(X, A \otimes Y^\sim) = \mathcal{Z}\left(\begin{array}{c} X \quad Y^\sim \\ \swarrow \quad \searrow \\ A \downarrow \end{array}\right).\end{aligned}$$

The orientation reversing morphism  $rev$  for internal arrow is obtained from the diagram of isomorphisms:

$$\begin{array}{ccc}\int^X \mathcal{Z}\left(\begin{array}{c} \quad \quad \quad \\ \swarrow \quad \searrow \\ \quad \quad \quad \end{array} \rightarrow X\right) \otimes \mathcal{Z}\left(\begin{array}{c} X \quad \quad \quad \\ \swarrow \quad \searrow \\ \quad \quad \quad \end{array}\right) & \xrightarrow{du_X \otimes du_X} & \int^{X \in \mathcal{C}} \mathcal{Z}\left(\begin{array}{c} \quad \quad \quad \\ \swarrow \quad \searrow \\ \quad \quad \quad \end{array} \rightarrow X\right) \otimes \mathcal{Z}\left(\begin{array}{c} X^\sim \quad \quad \quad \\ \swarrow \quad \searrow \\ \quad \quad \quad \end{array}\right) \\ \wr \downarrow & & \wr \downarrow \\ \mathcal{Z}\left(\begin{array}{c} \quad \quad \quad \\ \swarrow \quad \searrow \\ \quad \quad \quad \end{array} \rightarrow \quad \rightarrow \quad \leftarrow \quad \leftarrow \quad \begin{array}{c} \quad \quad \quad \\ \swarrow \quad \searrow \\ \quad \quad \quad \end{array}\right) & \xrightarrow{rev} \mathcal{Z}\left(\begin{array}{c} \quad \quad \quad \\ \swarrow \quad \searrow \\ \quad \quad \quad \end{array} \rightarrow \quad \rightarrow \quad \leftarrow \quad \leftarrow \quad \begin{array}{c} \quad \quad \quad \\ \swarrow \quad \searrow \\ \quad \quad \quad \end{array}\right) & \stackrel{f}{\simeq} \int^{Y \in \mathcal{C}} \mathcal{Z}\left(\begin{array}{c} \quad \quad \quad \\ \swarrow \quad \searrow \\ \quad \quad \quad \end{array} \rightarrow Y\right) \otimes \mathcal{Z}\left(\begin{array}{c} Y \quad \quad \quad \\ \swarrow \quad \searrow \\ \quad \quad \quad \end{array}\right)\end{array}$$

where the isomorphism  $f$  is that from Lemma 9.1. This shows that the relation (6.4) is satisfied. Similarly for (6.5), (6.6).

The existence of  $\mathcal{Z}(\text{rev}) : \mathcal{Z}(\Gamma) \rightarrow \mathcal{Z}(\Gamma')$  for arbitrary net  $\Gamma$  is guaranteed by

**Proposition 9.5.** *Let  $B : \mathcal{C}^{op} \times \mathcal{C} \rightarrow k\text{-vect}$  be a bifunctor. Then  $B_1(X, Y) = B(Y^\vee, X^\vee)$  is also a bifunctor. Let*

$$B(X, X) \xrightarrow{i_X} \int^X B(X, X), \quad B_1(Y, Y) \xrightarrow{j_Y} \int^Y B_1(Y, Y)$$

*be their coends. Then there exists the unique isomorphism  $\alpha$  of coends, which make the diagram*

$$\begin{array}{ccc} B(X^\vee, X^\vee) & \xrightarrow{i_{X^\vee}} & \int^X B(X, X) \\ \parallel & & \downarrow \alpha \\ B_1(X, X) & \xrightarrow{j_X} & \int^Y B_1(Y, Y) \end{array}$$

*commute for any  $X \in \mathcal{C}$ .*

**Proposition 9.6.** *The identity (6.7) is satisfied, that is, the diagram*

$$\begin{array}{ccccc} & & \text{Hom}(C, A \otimes B) & & \\ & d_r \nearrow & & \searrow d_l^{-1} & \\ \text{Hom}(A^\vee \otimes C, B) & & & & \text{Hom}(C \otimes B^\vee, A) \\ d_l^{-1} \uparrow & & & & \downarrow d_r \\ \text{Hom}(A^\vee, B \otimes C^\vee) & & & & \text{Hom}(B^\vee, C^\vee \otimes A) \\ & d_r \nwarrow & & \swarrow d_l^{-1} & \\ & & \text{Hom}(B^\vee \otimes A^\vee, C^\vee) & & \end{array}$$

*is commutative.*

The obtained functor satisfies (i)-(vii) and (xiii).

**9.5. Nets without ends.** We define also  $\mathcal{Z}(\Gamma)$  for a connected net without ends via insertion of a 1-vertex

$$\mathcal{Z}(\Gamma) \xrightarrow{\mathcal{Z}(\text{Ins})} \mathcal{Z}(\Gamma_\bullet) \simeq \mathcal{Z}(\Gamma_I).$$

Previous subsection shows that different choices give isomorphic answers. The property (xiii) is still true.

**9.6. Relations for fusing.** Now we construct a functor  $\mathcal{Z}$  on the category  $\mathcal{ON}$  extended by *ins*, *del*, *fus*. We put

$$\begin{aligned} \mathcal{Z}(fus) : \mathcal{Z} \left( \begin{array}{c} M \quad C \\ \swarrow \quad \searrow \\ \bullet \quad \bullet \\ \nearrow \quad \nwarrow \\ A \quad B \end{array} \right) &\simeq \int^{X \in \mathcal{C}} \text{Hom}(M, A \otimes X) \otimes \text{Hom}(X, B \otimes C) \simeq \\ &\simeq \text{Hom}(M, A \otimes (B \otimes C)) \xrightarrow{\text{Hom}(M, a_{A,B,C})} \text{Hom}(M, (A \otimes B) \otimes C) \simeq \\ &\simeq \int^{Y \in \mathcal{C}} \text{Hom}(M, Y \otimes C) \otimes \text{Hom}(Y, A \otimes B) \simeq \mathcal{Z} \left( \begin{array}{c} M \quad C \\ \swarrow \quad \searrow \\ \bullet \quad \bullet \\ \nearrow \quad \nwarrow \\ A \quad B \end{array} \right). \end{aligned}$$

The fusing pentagon (6.19) follows from the associativity pentagon in  $\mathcal{C}$ . We have to prove commutativity of diagrams (6.16), (6.17).

**Proposition 9.7.** *The relations (6.16), (6.17) are satisfied, that is, the diagrams*

$$\begin{array}{ccc} \text{Hom}(A, B \otimes (C \otimes D)) & \xrightarrow{\text{Hom}(A, a_{B,C,D})} & \text{Hom}(A, (B \otimes C) \otimes D) \\ \wr \downarrow & & \downarrow \wr \\ \int^X \text{Hom}(A, B \otimes X) \otimes \text{Hom}(X, C \otimes D) & & \int^X \text{Hom}(A, X \otimes D) \otimes \text{Hom}(X, B \otimes C) \\ \int 1 \otimes d_r^{-1} \downarrow & & \downarrow \int 1 \otimes d_l^{-1} \\ \int^X \text{Hom}(A, B \otimes X) \otimes \text{Hom}(C^\sim \otimes X, D) & & \int^X \text{Hom}(A, X \otimes D) \otimes \text{Hom}(X \otimes C^\sim, B) \\ \int d_l^{-1} \otimes d_l \downarrow & & \downarrow \int d_r^{-1} \otimes d_r \\ \int^X \text{Hom}(A \otimes X^\sim, B) \otimes \text{Hom}(C^\sim, D \otimes X^\sim) & & \int^X \text{Hom}(X^\sim \otimes A, D) \otimes \text{Hom}(C^\sim, X^\sim \otimes B) \\ \wr \downarrow & & \downarrow \wr \\ \int^Y \text{Hom}(A \otimes Y, B) \otimes \text{Hom}(C^\sim, D \otimes Y) & & \int^Y \text{Hom}(Y \otimes A, D) \otimes \text{Hom}(C^\sim, Y \otimes B) \\ \int d_r \otimes 1 \downarrow & & \downarrow \int d_l \otimes 1 \\ \int^Y \text{Hom}(Y, A^\sim \otimes B) \otimes \text{Hom}(C^\sim, D \otimes Y) & & \int^Y \text{Hom}(Y, D \otimes A^\sim) \otimes \text{Hom}(C^\sim, Y \otimes B) \\ \wr \downarrow & & \downarrow \wr \\ \text{Hom}(C^\sim, D \otimes (A^\sim \otimes B)) & \xrightarrow{\text{Hom}(C^\sim, a_{D,A^\sim,B})} & \text{Hom}(C^\sim, (D \otimes A^\sim) \otimes B) \end{array}$$

$$\begin{array}{ccc}
\mathrm{Hom}(D, A \otimes (B \otimes C)) & \xrightarrow{\mathrm{Hom}(D, a_{A,B,C},)} & \mathrm{Hom}(D, (A \otimes B) \otimes C) \\
\uparrow \wr & & \downarrow \wr \\
\int^X \mathrm{Hom}(D, A \otimes X) \otimes \mathrm{Hom}(X, B \otimes C) & & \int^X \mathrm{Hom}(D, X \otimes C) \otimes \mathrm{Hom}(X, A \otimes B) \\
\uparrow \int d_r \otimes 1 & & \downarrow \int 1 \otimes d_r^{-1} \\
\int^X \mathrm{Hom}(A^\vee \otimes D, X) \otimes \mathrm{Hom}(X, B \otimes C) & & \int^X \mathrm{Hom}(D, X \otimes C) \otimes \mathrm{Hom}(A^\vee \otimes X, B) \\
\uparrow \int d_l^{-1} \otimes 1 & & \downarrow \int d_r^{-1} \otimes d_l \\
\int^X \mathrm{Hom}(A^\vee, X \otimes D^\vee) \otimes \mathrm{Hom}(X, B \otimes C) & & \int^X \mathrm{Hom}(X^\vee \otimes D, C) \otimes \mathrm{Hom}(A^\vee, B \otimes X^\vee) \\
\uparrow \wr & & \downarrow \wr \\
\mathrm{Hom}(A^\vee, (B \otimes C) \otimes D^\vee) & & \int^Y \mathrm{Hom}(Y \otimes D, C) \otimes \mathrm{Hom}(A^\vee, B \otimes Y) \\
\uparrow \mathrm{Hom}(A^\vee, a_{B,C,D^\vee}) & & \downarrow \int d_l \otimes 1 \\
\mathrm{Hom}(A^\vee, B \otimes (C \otimes D^\vee)) & \xleftarrow{\sim} & \int^Y \mathrm{Hom}(Y, C \otimes D^\vee) \otimes \mathrm{Hom}(A^\vee, B \otimes Y)
\end{array}$$

are commutative.

**9.7. Relations for braiding and twists.** We extend the functor  $\mathcal{Z}$  to the category generated over  $\mathcal{ON}$  by *ins*, *del*, *fus*, *Tw* and *Br*. We prove the relations involving *Tw* and *Br*. All conditions with *Tw* are obvious or follow from equations

$$\nu \begin{array}{|c|} \hline \square \\ \hline \end{array} \begin{array}{|c|} \hline \cup \\ \hline \end{array} = \begin{array}{|c|} \hline \cup \\ \hline \end{array} \nu \begin{array}{|c|} \hline \square \\ \hline \end{array}, \quad \nu \begin{array}{|c|} \hline \square \\ \hline \end{array} \begin{array}{|c|} \hline \cup \\ \hline \end{array} = \begin{array}{|c|} \hline \cup \\ \hline \end{array} \nu \begin{array}{|c|} \hline \square \\ \hline \end{array}.$$

The hexagon (6.25) for braiding

$$\mathrm{Hom}(C, A \otimes B) \xrightarrow{\mathrm{Hom}(C, c_{AB})} \mathrm{Hom}(C, B \otimes A)$$

follows from that one for commutativity  $c$ . Property (6.22) follows from the equation

$$\mathrm{Hom}(C, c_{AB}^2) = \mathrm{Hom}(C, \nu_{A \otimes B} \circ \nu_A^{-1} \otimes \nu_B^{-1}) = \mathrm{Hom}(\nu_C, \nu_A^{-1} \otimes \nu_B^{-1}).$$

**Proposition 9.8.** *The relations (6.23), (6.24) are satisfied. That is, the diagrams*

$$\begin{array}{ccccc}
\text{Hom}(C, A \otimes B) & \xrightarrow{\text{Hom}(C, c)} & \text{Hom}(C, B \otimes A) & \xrightarrow{d_r^{-1}} & \text{Hom}(B^\sim \otimes C, A) \\
\uparrow & & & & \downarrow d_l \\
\text{Hom}(C, \nu \otimes B) & & & & \text{Hom}(B^\sim, A \otimes C^\sim) \\
& & & & \downarrow \text{Hom}(B^\sim, c) \\
\text{Hom}(C, A \otimes B) & \xleftarrow{d_l} & \text{Hom}(C \otimes B, A) & \xleftarrow{d_r^{-1}} & \text{Hom}(B^\sim, C^\sim \otimes A) \\
\\ 
\text{Hom}(C, A \otimes B) & \xrightarrow{\text{Hom}(C, c)} & \text{Hom}(C, B \otimes A) & \xrightarrow{d_l^{-1}} & \text{Hom}(C \otimes A^\sim, B) \\
\uparrow & & & & \downarrow d_r \\
\text{Hom}(C, A \otimes \nu) & & & & \text{Hom}(A^\sim, C^\sim \otimes B) \\
& & & & \downarrow \text{Hom}(A^\sim, c) \\
\text{Hom}(C, A \otimes B) & \xleftarrow{d_r} & \text{Hom}(A^\sim \otimes C, B) & \xleftarrow{d_l^{-1}} & \text{Hom}(A^\sim, B \otimes C^\sim)
\end{array}$$

*commute.*

**9.8. A quotient functor.** Let  $g : \Gamma' \rightarrow \Gamma$  be a glueing and  $gf_1 = gf_2 : \Gamma' \rightarrow \Gamma''$ . Then  $\mathcal{Z}(g)\mathcal{Z}(f_1) = \mathcal{Z}(g)\mathcal{Z}(f_2)$ . The morphism  $\mathcal{Z}(g) : \mathcal{Z}(\Gamma'; A, X; X, B) \rightarrow \mathcal{Z}(\Gamma; A; B)$  can be regarded as a morphisms of left exact functors belonging to  $\hat{\mathcal{C}}_{k,l}$

$$\mathcal{Z}(g) : \oplus_{X \in \mathcal{C}^n} \mathcal{Z}(\Gamma'; A, X; X, B) \longrightarrow \mathcal{Z}(\Gamma; A; B).$$

It is an epimorphism if  $k + l > 0$ . Hence, by definition  $\mathcal{Z}(f_1) = \mathcal{Z}(f_2)$ .

In case  $k = l = 0$  we add one more leg  $A$  to our nets, obtaining new morphisms  $\bar{g}\bar{f}_1 = \bar{g}\bar{f}_2$ . By the above considerations  $\mathcal{Z}(\bar{f}_1) = \mathcal{Z}(\bar{f}_2) : \mathcal{Z}(\bar{\Gamma}; A; ) \rightarrow \mathcal{Z}(\bar{\Gamma}''; A; )$ . Setting  $A = I$  and applying isomorphisms we change the end  $A$  to a 1-vertex. Deleting it we again deduce  $\mathcal{Z}(f_1) = \mathcal{Z}(f_2)$ .

When  $g$  is any other generator it is invertible, hence, in all cases we proved that  $gf_1 = gf_2$  implies  $\mathcal{Z}(f_1) = \mathcal{Z}(f_2)$ . This means that the functor  $\mathcal{Z}$  is, in fact, defined on a category having the left cancellation property.

Now we construct a functor  $Z$  as an epimorphic image of  $\mathcal{Z}$  on the category with left cancellations  $\mathcal{ON}$  extended by *ins*, *del*, *fus*, *Tw*, *Br* to  $k$ -vect.

**Proposition 9.9.** *There exists a unique up to equivalence functor  $Z$  on the category  $\mathcal{ON}$  extended by *ins*, *del*, *fus*, *Tw*, *Br* to  $k$ -vect satisfying*



conditions (i)–(xi) of Section 9.2. It has also the property (xii). There is an epimorphism  $\mathcal{Z} \rightarrow \mathcal{Z}$ .

**9.9. Construction of switches.** Now we extend the functor  $Z$  to  $EN$ . We shall show that there are exactly two such extensions, which differ by a sign of  $Z(S)$ .

Morphisms  $S^{\pm 1}$  in  $EN$  must be represented by functorial in  $X$  isomorphisms

$$\mathrm{Hom}(X, \mathbf{f}) \simeq Z\left(\begin{array}{c} X \\ \xrightarrow{\quad} \bullet \end{array} \begin{array}{c} \circlearrowright \end{array}\right) \xrightarrow{Z(S^{\pm 1})} Z\left(\begin{array}{c} X \\ \xrightarrow{\quad} \bullet \end{array} \begin{array}{c} \circlearrowright \end{array}\right) \simeq \mathrm{Hom}(X, \mathbf{f}).$$

Hence, they are induced by automorphisms  $S^{\pm 1} : \mathbf{f} \rightarrow \mathbf{f}$  in  $\mathcal{C}$ .

**Proposition 9.10.** *The axiom (6.28) is satisfied, or equivalently the diagram at Figure 7 made of morphisms of left exact functors in  $X, Y$  (quotients are taken in the category of left exact functors) with  $K = \text{Ker}(\int^M M \otimes M^\sim \rightarrow \mathbf{f})$  is commutative if and only if*

$$S^{-1} = \text{diagram} \quad (9.1)$$

for some morphism  $\mu : I \rightarrow \mathbf{f}$ .

The relation (6.27) implies that

$$\begin{array}{ccccc} \int^X X \otimes X^\sim & \xrightarrow{f^c} & \int^X X^\sim \otimes X & \xrightarrow{f^{1 \otimes \nu}} & \int^X X^\sim \otimes X \\ i_X \downarrow & & & & \downarrow i_{X^\sim} \\ \mathbf{f} & \xrightarrow{S^{-2}} & & & \mathbf{f} \end{array}$$

by Proposition 9.5. That is,

$$S^{-2} = \gamma : \mathbf{f} \rightarrow \mathbf{f} \quad (9.2)$$

$$\begin{array}{ccc}
\mathrm{Hom}(Y \otimes X, \mathbf{f}) & \xrightarrow{\mathrm{Hom}(c_{XY}, \mathbf{f})} & \mathrm{Hom}(X \otimes Y, \mathbf{f}) \\
\mathrm{Hom}(Y \otimes X, S^{-1}) \downarrow & & \downarrow \wr \\
\mathrm{Hom}(Y \otimes X, \mathbf{f}) & & \mathrm{Hom}(X \otimes Y, \int^N N \otimes N^\vee) / \mathrm{Hom}(X \otimes Y, K) \\
\wr \downarrow & & \downarrow \wr \\
\frac{\mathrm{Hom}(Y \otimes X, \int^M M \otimes M^\vee)}{\mathrm{Hom}(Y \otimes X, K)} & & \oint^N \mathrm{Hom}((X \otimes Y) \otimes N, N) / \mathrm{Hom}(X \otimes Y, K) \\
\wr \downarrow & & \downarrow \mathrm{Hom}(a, N) \\
\oint^M \frac{\mathrm{Hom}((Y \otimes X) \otimes M, M)}{\mathrm{Hom}(Y \otimes X, K)} & & \oint^N \mathrm{Hom}(X \otimes (Y \otimes N), N) / \mathrm{Hom}(X \otimes Y, K) \\
\mathrm{Hom}(a, M) \downarrow & & \downarrow \wr \\
\oint^M \frac{\mathrm{Hom}(Y \otimes (X \otimes M), M)}{\mathrm{Hom}(Y \otimes X, K)} & & \oint^{N, P} \frac{\mathrm{Hom}(X \otimes P, N) \otimes \mathrm{Hom}(Y \otimes N, P)}{\mathrm{Hom}(X \otimes Y, K)} \\
\mathrm{Hom}(Y \otimes \nu_{X \otimes M}^{-1}, \nu) \downarrow & & \downarrow \wr \\
\oint^M \frac{\mathrm{Hom}(Y \otimes (X \otimes M), M)}{\mathrm{Hom}(Y \otimes X, K)} & & \oint^P \mathrm{Hom}(Y \otimes (X \otimes P), P) / \mathrm{Hom}(X \otimes Y, K) \\
\mathrm{Hom}(a^{-1}, M) \downarrow & & \downarrow \mathrm{Hom}(a^{-1}, P) \\
\oint^M \frac{\mathrm{Hom}((Y \otimes X) \otimes M, M)}{\mathrm{Hom}(Y \otimes X, K)} & & \oint^P \mathrm{Hom}((Y \otimes X) \otimes P, P) / \mathrm{Hom}(Y \otimes X, K) \\
\wr \downarrow & & \downarrow \wr \\
\frac{\mathrm{Hom}(Y \otimes X, \int^M M \otimes M^\vee)}{\mathrm{Hom}(Y \otimes X, K)} & & \mathrm{Hom}(Y \otimes X, \int^P P \otimes P^\vee) / \mathrm{Hom}(Y \otimes X, K) \\
\wr \downarrow & & \downarrow \wr \\
\mathrm{Hom}(Y \otimes X, \mathbf{f}) & \xrightarrow{\mathrm{Hom}(Y \otimes X, S)} & \mathrm{Hom}(Y \otimes X, \mathbf{f})
\end{array}$$

FIGURE 7. The realization of a relation for a torus with two holes

(see Section 8). So we find

$$S = \gamma^{-1} S^{-1} =$$

$$(9.3)$$

Theorem 6.13 from [10] states that if morphisms (9.1) and (9.3) are inverse to each other, then  $\mu$  is an integral of the Hopf algebra  $\mathbf{f}$ . It is unique up to a constant. The normalizing constant for the integral  $\mu$  is fixed up to a sign by (9.2). The relation

$$(ST)^3 = \lambda S^2$$

for  $T = \int 1 \otimes \nu : \mathbf{f} \rightarrow \mathbf{f}$  and some constant  $\lambda$  is proven in [10]. Hence, putting the central charge  $C$  equal this constant  $\lambda$  on  $Z(A_1)$  we get the relation (7.5). For arbitrary net  $\Gamma$  of genus  $g$  we set  $Z(C_\Gamma) = \lambda^g$ .

Thus, we obtained a functor  $EN \rightarrow k\text{-vect}$  satisfying all conditions of Section 9.2. It is unique up to a choice of sign of the normalizing constant.

## REFERENCES

1. J. S. Birman, *Mapping class groups and their relationship to braid groups*, Commun. Pure Appl. Math. **22** (1969), 213–238.
2. M. Dehn, *Die Gruppe der Abbildungsklassen*, Acta Math. **69** (1938), 135–206.
3. A. Grothendieck, *Esquisse d'un programme*, 1984, unpublished.
4. A. Joyal and R. Street, *Tortile Yang-Baxter operators in tensor categories*, J. Pure Appl. Alg. **71** (1991) 43–51.
5. T. Kohno, *Topological invariants for 3-manifolds using representations of mapping class groups I*, Topology **31** (1992), n. 2, 203–230.
6. ———, *Three-manifold invariants derived from conformal field theory and projective representations of modular groups*, Int. J. of Modern Phys. B **6** (1992), n. 11–12, 1795–1805.
7. W. B. R. Lickorish, *A representation of orientable combinatorial 3-manifolds*, Annals Math. **76** (1962), n. 3, 531–540.
8. ———, *A finite set of generators for the homeotopy group of a 2-manifold*, Proc. Camb. Phil. Soc. **60** (1964), 769–778.
9. V. V. Lyubashenko, *Tangles and Hopf algebras in braided categories*, to appear in J. Pure Appl. Alg.
10. ———, *Modular transformations for tensor categories*, to appear in J. Pure Appl. Alg.
11. ———, *Ribbon categories as modular categories*, preprint.
12. V. V. Lyubashenko and S. Majid, *Braided groups and quantum Fourier transform*, to appear in J. Algebra.

13. S. Mac Lane, *Categories for the working mathematician*, Springer-Verlag, 1971.
14. W. Magnus, *Über Automorphismen von Fundamentalgruppen berandeter Flächen*, Math. Ann. **109** (1934), 617–646.
15. S. Majid, *Braided groups*, Preprint, DAMTP/90-42, 1990.
16. G. Moore and N. Seiberg, *Classical and Quantum Conformal Field Theory*, Commun. Math. Phys., **123** (1989), 177–254.
17. N. Reshetikhin, *Quasitriangular Hopf algebras, solutions to the Yang-Baxter equations and link invariants*, Algebra and Analysis **1** (1989), n. 2, 169–194.
18. N. Reshetikhin and V. G. Turaev, *Invariants of 3-manifolds via link polynomials and quantum groups*, Invent. math. **103** (1991), 547–597.
19. G. P. Scott, *Braid groups and the group of homeomorphisms of a surface*, Proc. Camb. Phil. Soc. **68** (1970), 605–617.
20. M. C. Shum, *Tortile tensor categories*, Macquarie preprint, 1989.
21. V. Turaev, *Quantum invariants of 3-manifolds*, Publ. de l'IRMA 509/P-295, October 1992.
22. B. Wajnryb, *A simple presentation for the mapping class group of an orientable surface*, Israel J. Math. **45** (1983), no. 2–3, 157–174.
23. K. Walker, *On Witten's 3-manifold invariants*, Preprint, February 1991.
24. E. Witten, *Quantum field theory and the Jones polynomial*, Commun. Math. Phys. **121** (1989), 351–399.